

REVIEW

Open Access



Tension at the gate: sensing mechanical forces at the blood–brain barrier in health and disease

Cathrin E. Hansen^{1,2,3†}, David Hollaus^{1†}, Alwin Kamermans^{1,2} and Helga E. de Vries^{1,2,3*}

Abstract

Microvascular brain endothelial cells tightly limit the entry of blood components and peripheral cells into the brain by forming the blood–brain barrier (BBB). The BBB is regulated by a cascade of mechanical and chemical signals including shear stress and elasticity of the adjacent endothelial basement membrane (BM). During physiological aging, but especially in neurological diseases including multiple sclerosis (MS), stroke, small vessel disease, and Alzheimer's disease (AD), the BBB is exposed to inflammation, rigidity changes of the BM, and disturbed cerebral blood flow (CBF). These altered forces lead to increased vascular permeability, reduced endothelial reactivity to vasoactive mediators, and promote leukocyte transmigration. Whereas the molecular players involved in leukocyte infiltration have been described in detail, the importance of mechanical signalling throughout this process has only recently been recognized. Here, we review relevant features of mechanical forces acting on the BBB under healthy and pathological conditions, as well as the endothelial mechanosensory elements detecting and responding to altered forces. We demonstrate the underlying complexity by focussing on the family of transient receptor potential (TRP) ion channels. A better understanding of these processes will provide insights into the pathogenesis of several neurological disorders and new potential leads for treatment.

Keywords Blood–brain barrier, Mechanical forces, Wall shear stress, Basement membrane stiffness, Transendothelial migration, TRP, Multiple sclerosis, Alzheimer's disease, Stroke, Small vessel disease

Background

Exchanges of circulating molecules and cells between the bloodstream and central nervous system (CNS) are regulated by the neurovascular unit, a multi-layered structure

consisting of perivascular astrocytes, microglia, pericytes, and specialized brain microvascular endothelial cells (BMECs) comprising the blood–brain barrier (BBB) [1–3] (Fig. 1, central part). The BBB is the innermost, monocellular layer of the brain microvasculature forming a tight physical barrier that is characterized by a unique network of adherens junctions (AJs) and tight junctions (TJs) [3, 4], which respond to internal and external mechanical forces of the local microenvironment [5, 6] (Fig. 1B). AJs consist of transmembrane protein complexes, such as vascular endothelial (VE-) cadherin [1, 4], which connect to the intracellular actin cytoskeleton via catenins [4, 7]. TJs are located at the basolateral side of BMECs and include claudin-5, occludin, junctional adhesion molecules (JAMs), and tri-cellular junction proteins

[†]Cathrin E. Hansen and David Hollaus have contributed equally to this work.

*Correspondence:

Helga E. de Vries
he.devries@amsterdamumc.nl

¹ Department of Molecular Cell Biology and Immunology, Amsterdam UMC location Vrije Universiteit Amsterdam, De Boelelaan 1117, Amsterdam, The Netherlands

² Amsterdam Neuroscience, Amsterdam UMC, Amsterdam, The Netherlands

³ MS Center Amsterdam, Amsterdam UMC Location VU Medical Center, Amsterdam, The Netherlands



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.




Fig. 1 A trio of mechanical forces acting on the brain endothelium. Forces and their directionality are presented by black arrows. **A** Wall shear stress (WSS) acts on the luminal side of the BMECs in the direction of the cerebral blood flow (CBF). **B** Abluminally, BMECs are anchored to the basement membrane (BM) and use integrin-mediated mechanotransduction to respond to the underlying stiffness. Basolateral junction complexes include tight junctions (TJs) and adherens junctions (AJs) creating inter- and intracellular tension. **(C)** Mostly under pathological conditions: Immune cell migration across the BBB is a multi-step cascade with mechanical forces acting on both cellular players, the immune cell, and BMECs. Figure is based on Vanlandewijk et al. [37]. *HSPGs* Heparan sulfate proteoglycans, *Coll IV* Collagen IV, *PVS* perivascular space, *BM* basement membrane

such as tricellulin [8–10]. Together, they form a robust paracellular connection and generate intercellular tension, giving rise to the exceptionally low permeability of the BBB [1, 3]. Finally, integrin-based focal adhesions anchor BMECs to the underlying basement membrane (BM), a network of extracellular matrix (ECM) proteins, and an important source of mechanical force (Fig. 1B) [11]. To maintain their integrity, BMECs sense and adapt to changes in forces including cerebral blood flow (CBF), BM elasticity, and cell–cell interactions such as during transendothelial migration (TEM) of peripheral immune cells across the BBB (Fig. 1A–C) [1, 4, 8].

Pathological changes of the BBB are found in many cerebral disorders. In multiple sclerosis (MS), a chronic neuroinflammatory disease, BBB impairment, and immune cell infiltration are disease hallmarks and tightly connected to the formation of demyelinated plaques [12, 13]. Alzheimer's disease (AD) is characterized by aggregation of amyloid beta (A β) and

accumulation of hyperphosphorylated tau forming neurofibrillary tangles [14]. These are further accentuated by vascular changes including BBB injury, reduced CBF, and enlarged perivascular spaces [15]. Impaired A β clearance is a common feature of AD and cerebral amyloid angiopathy (CAA), whereas in the latter A β accumulates particularly around the vasculature, further impairing vascular reactivity and possibly permitting haemorrhages [16]. Reduced vascular function through vessel stiffening and chronic hypertension is commonly found in aging cerebral vasculature and is known to increase the risk for stroke (haemorrhagic and ischemic) and sporadic small vessel disease [17, 18]. Overall, vascular stiffening and disturbed CBF likely result in reduced vascular reactivity, enhanced perivascular aggregate formation, and eventually focal hypoxia, contributing to the (vascular) pathology observed in numerous age-related neurodegenerative disorders [19–28].

Many detrimental vascular changes in cerebral disorders follow and include distinct mechanical components such as alterations of BM composition and wall shear stress (WSS) acting on the microvasculature (Fig. 1A, B). WSS or often referred to as simply shear stress defines the arising force within a moving fluid and between the fluid and vessel wall [29]. Additionally, neuroinflammation facilitates the migration of peripheral immune cells across the BBB, which adds cell–cell interaction forces to the mechanical trio [30, 31] (Fig. 1C). TEM of immune cells into the brain is a hallmark of MS, but also gains increasing attention in AD patients, stroke, and healthy elderly [32, 33]. This migration cascade governs many biomechanical and biochemical interactions from start to finish. Whereas biochemical signalling has been described and reviewed in detail [1, 3, 8, 34], the importance of mechanical stress has only recently been realised.

It becomes apparent that changes in mechanical forces largely influence vascular function and dysfunction and might be early drivers of neuroinflammatory and neurodegenerative diseases. In this review, we briefly summarize our current knowledge of the two main mechanical stressors, WSS and BM stiffness, acting on the BBB and their pathological changes. Then, we highlight the forces measured during TEM of immune cells across the BBB and BBB-specific migration behaviour. Finally, we focus on mechanosensory/sensitive elements of BMECs and spotlight members of the transient receptor potential (TRP) ion channel family to showcase the intricacies of mechanical signalling [35, 36].

Mechanical forces at the BBB

Cerebral vessels are subjected to a range of mechanical stressors (Table 1), which play a key role in BBB maintenance, activation, and function [38, 39]. Changes in mechanical stimuli are sensed by specialized mechanosensors and translated into biochemical signalling cascades [40], culminating in cytoskeletal reorganization, gene regulation, and epigenetic chromatin modification [41]. Disturbances in mechanical forces including CBF and BM stiffness are often concomitant and recognized as early signs of numerous cerebral disorders [5, 39]. Below, the force ranges and impact on the cerebral vasculature are discussed.

External, non-cellular forces

Wall shear stress

Wall shear stress (WSS) or endothelial WSS is calculated from the blood viscosity (highly dependent on concentration and mechanical properties of red blood cells) and axial flow velocity gradient at the vessel wall [66]. WSS, arising from laminar or disturbed CBF at the luminal side of BMECs, plays a critical role in maintaining the structure and function of the BBB [38]. Under healthy conditions, the average capillary WSS in the brain ranges from 1 to 2 N/m² [46, 47] (Table 1, Fig. 1A). ECs experience the highest WSS within capillaries and declining hyperbolically to 0.28 N/m² at larger diameter post-capillary venules [48, 67]. This WSS was shown to be required for the differentiation of vascular ECs into a BMEC phenotype [68] using a dynamic in vitro BBB model. In vitro application of capillary-like WSS also inhibited cell proliferation, probably due to induction of the cyclin-dependent kinase inhibitor p21, causing cell cycle arrest [5, 69]. Recent advances in BBB models, which incorporate external

Table 1 Mechanical stressors and forces acting at the vasculature. Summary of non-cellular and cellular mechanical forces at the endothelium measured in N/m² (equals 10 dyne/cm² or 1 Pa)

Stressor	Area/cell type	Force range	References
External non-cellular forces			
Wall shear stress	Arteries	0.4–3 N/m ²	[39, 42–45]
	Arterioles & capillaries	Avg. 1–2 N/m ²	[46–50]
	Post-capillary venules & veins	< 0.1–4 N/m ²	[39, 48, 49, 51]
BM stiffness	Arteries & microvessels	~ 50 K to 150 K N/m ² or higher	[39, 52–55]
	Venules/venous tissue (outside brain)	~ 3 K–50 K N/m ²	[54, 56]
Cell–cell interaction forces			
Selectin catch bonds	Leukocytes & ECs	10–30 pN/bond, maybe up to 60 pN	[57–60]
Integrin catch bonds	Leukocytes & ECs	10–30 pN/bond	[61, 62]
Crawling	T cells	up to 0.6 nN/cell	[63]
Diapedesis	Leukocytes	up to 60 nN/cell	[58]
Cell–cell junctions e.g. VE-cadherin and PECAM-1 binding	ECs	10–100 pN/bond (depending on number and type of interacting proteins)	[64, 65]

mechanical forces such as shear stress and ECM stiffness, have been reviewed elsewhere [70–74].

CBF dynamics strongly alter with age, occlusion and inflammation, whereas both high and low WSS are associated with brain pathology [75]. Aberrant high WSS (>4 N/m²) occurs in hypertension and causes vascular dysfunction through loss of TJ proteins and activation of inflammatory signalling at the BBB [38, 76–78]. Hypertension and cerebrovascular injury can manifest in small vessel disease, which often presents an intermediate state leading to ~30% of ischemic stroke and ~80% of haemorrhagic intracerebral bleedings [79–82]. In small vessel disease, the stiffened/damaged arteries are unfit to absorb the heightened pulsatility, causing damage to the proximate microvasculature of the brain [83, 84]. In vitro studies confirm that pulsatile WSS on BMECs causes the loss of TJ proteins, reduced P-glycoprotein expression and delocalization of zona occludens-1 (ZO-1) from the cell borders to the cytoplasm and nucleus [77]. Disease severity of small vessel disease is also associated with global and local CBF alterations and recent work has pinpointed distinct region-specific CBF patterns in patients, which were linked to the occurrence of periventricular white matter hyperintensities and total disease burden [85].

During ischemic stroke, blood flow is blocked to a part of the cerebral vascular network, instantly severely lowering CBF in this region, which can be restored upon timely treatment (reperfusion). As a result, BMECs first experience hypoxia and oxidative stress which partially increases their permeability for blood components, fluids, and peripheral cells, causing post-ischemic inflammation [86]. Edema formation follows ischemia-induced sodium and water uptake through shear stress-regulated ion transporter [87]. A number of signalling pathways (including Wnt and Notch pathways) underlying ischemic/reperfusion injury across organs have been recently reviewed [88]. Very low levels of CBF and WSS also prompt increased BMEC proliferation and enhanced BBB permeability through cell layer remodelling [38, 69, 89]. Following reperfusion, BMECs secondly experience enhanced WSS which promotes endothelial-to-mesenchymal transition [90]. This (partial) shift to mesenchymal cell properties further enhances BBB permeability [91], which may however reverse at a later timepoint of reperfusion [91]. A recent study, using two-photon imaging to follow vascular leakage post ischemic stroke in mice, showed BBB leakage as early as 30 min after the occlusion, steadily increasing in the hours after (120 min) [92]. Disruption of BBB function from ischemia/reperfusion has been reported as a predictor of poor disease outcome and haemorrhagic transformation in ischemic stroke patients [93].

Hypertension-induced vascular injury correlates with age and increases the risk for AD, yet the CBF is frequently reduced globally and locally in AD patients, which has been suggested as an early marker for AD [20, 94–96]. Similarly, MS patients show reduced CBF in white matter regions, specifically periventricular areas, concomitant with increased BBB permeability [97, 98]. Reduced CBF in MS may be the result of impaired flow-mediated endothelial dilation, both of which are correlated with disease severity and progression [99]. Whereas numerous studies hint towards changes in CBF in age-related and neuroinflammatory diseases [100, 101], the underlying causes and consequences for BMECs and BBB function remain elusive. Especially data on human brain vasculature is missing to fully comprehend different forms of WSS and its effects on the BBB during aging and neuroinflammatory disorders [51].

Basement membrane changes

Vascular stiffness is a complex property governed chiefly by vascular reactivity, BM properties, and blood pressure [102]. Here we focus on the BM and its changes in mechanical properties during pathology. The brain endothelial BM separates BMECs from neighbouring pericytes and supports BBB maintenance [103]. The BM is composed of extracellular matrix (ECM) proteins including laminins, nidogens, heparan sulphate proteoglycans, and collagen IV (Fig. 1B). They form a 50–100 nm thick three-dimensional network that interacts with BMECs to hold them in place [104, 105]. BM stiffness is influenced by the composition of ECM proteins, their concentration and elasticity, crosslinking between the proteins and the created intrinsic tension [106]. Degradation through proteases (e.g. matrix metalloproteases), remodelling and (limited) synthesis of ECM proteins can change the degree of membrane stiffness [106].

In vitro studies show that ECs respond to the rigidity of the neighbouring BM by enhanced expression of ECM receptors, including integrins. These cell–matrix contacts are crucial for BMEC identity and BBB function [107, 108]. Consequently, global knockout models for components of the ECM or pharmacological blockage of integrins often demonstrate cerebral bleeds and pre- or postnatal lethality [109, 110]. Growing cells on polyacrylamide hydrogels of varying stiffness showed that cells on soft hydrogels developed a lower density of actin fibres and more rounded nuclei than those grown on stiffer substrates [39, 111]. BM stiffness further cues endothelial progenitor cell differentiation along the arterial-venous axis, as higher stiffness induces arterial EC marker expression via the Ras-Mek pathway [112]. Interestingly, when exposed to physiological WSS, ECs on soft hydrogels displayed tighter cell junctions, more profound

elongation, and fewer (RhoA-mediated) cytoskeletal changes, revealing a direct relation between ECM stiffness and WSS [111]. These results show that ECs can directly modulate internal tension in response to external forces by altering the assembly and dynamics of actin fibres and junctional proteins [40, 113].

BM thickening has been linked to a narrowing of the vessel lumen during inflammation, aging and AD [114–118]. Over time, the BM composition shifts towards thicker collagen fibres with enhanced crosslinking, while hyaluronan levels decrease by degradation and elastin elasticity function is reduced [119, 120]. Consequent narrowing of the vessel lumen diameter restricts local CBF, causing lack of sufficient oxygen and nutrient supply for the brain [121, 122]. Prolonged reduced CBF can be further amplified by decreased endothelial-dependent dilation and overactive contraction, e.g. by pericytes evoked through reactive oxygen species or A β signalling in ischemic stroke and AD [123, 124]. Moreover, the increase in BM rigidity affects mechanosensitive processes such as angiogenesis [125, 126]. Functional angiogenesis requires a delicate balance between matrix remodelling/degradation and mechanical support of the BM for BMEC adhesion and migration. Age-related changes in BM composition and elasticity result in a lack of vessel support and might be linked to microvessel rarefaction and regression following impaired angiogenesis [127]. Lastly, the endothelial BM is a reservoir for growth factors and signalling crucial for vessel formation. Changing ECM protein composition thus also leads to a dysregulation in growth factor and other ligand availability [127].

In AD pathogenesis, changes in levels of collagen IV and the proteoglycan perlecan occur early on (Braak stage > 2 & $\leq$ 4) and correlate with parenchymal A β plaque deposition in the frontal and temporal cortex, suggesting BM changes in parallel with disease progression [128]. In AD patients with CAA, CAA severity correlates with A β _{1–42} and collagen IV abundance in vessels of the frontal cortex [129]. It remains to be investigated if changes in the endothelial BM facilitate vascular A β deposition or merely occur concomitant with disease development. In MS, non-invasive pulse wave velocity measurements, a readout for arterial stiffness, are associated with disease duration and severity. Another study showed that increased arterial stiffness correlated with a reduced cognitive processing speed, as assessed via the Symbol Digit Modalities Test [130, 131]. Similar measurements in stroke patients connected arterial stiffness with enlarged perivascular spaces and cerebral microbleeds and revealed vessel stiffness as a potentially more important risk factor than blood pressure for cerebral small vessel disease [132, 133]. Lastly, a meta-analysis demonstrated

a negative association between arterial stiffness and cognitive function, together highlighting the importance of vascular fitness in disease pathogenesis and global memory function [134, 135].

Due to their dense network, BMs also create a rate-limiting step for immune cells to enter the perivascular space and the parenchyma, prolonging the migration duration by 3–fourfold [136–138]. Pathological changes in BM composition can facilitate the infiltration of immune cells and cue cell differentiation as previously reviewed elsewhere [139]. Matrikines, peptides released during ECM remodelling, might also affect BMEC phenotype and inflammatory state as suggested for the peripheral endothelium [140, 141]. Ex vivo measurements of BMs have shown a large range of stiffness (Young's elastic module) from 500 to 4,000,000 N/m² depending on the tissue, but similar to WSS, measuring the stiffness of BMs in vivo remains an ongoing challenge (Table 1) [142].

Cell–cell interaction forces during transendothelial migration

Immune cell migration across the BBB under homeostasis is limited to a small subset of activated immune cells [143]. Dysfunction of the BBB due to inflammatory and age-related pathology greatly enhances TEM. Immune cell entry further potentiates inflammatory processes within the CNS, fuelling disease progression and cognitive decline [144–146]. In addition to chemical cues, leukocytes prepare for migration by following a stiffness gradient of the underlying substrate, a process termed durotaxis [147, 148]. In the following sections, we highlight the forces acting between BMECs and immune cells during the multi-step process of TEM. Studies on the mechanical forces between BMECs and immune cells have predominantly focused on T cells, which are key players in neuropathological events. However, many of these mechanisms are likely universal to immune cell populations entering the CNS.

First steps of engagement of leukocytes with BMECs

In order to exit the bloodstream and enter the brain parenchyma, immune cells must come into contact with the vessel wall [149]. Thus, migration primarily takes place at inflamed postcapillary venules where WSS is greatly reduced, increasing the chance of immune cells interacting with the brain endothelium [150]. This is further facilitated by erythrocytes, which are thought to push immune cells to the edges of the blood vessel [149] (Fig. 2).

The first contacts of the leukocyte with the endothelial cells (ECs) are generally made between the family of molecules known as selectins (L-selectin on leukocytes, P- and E-selectin on ECs) and/or α_4 -integrins (such as




Fig. 2 TEM cascade of migratory T cells across the brain endothelium. Upon engagement of catch bonds between selectins/ α_4 -integrins and their ligands, T cells start rolling and eventually firmly adhere to the ECs. Cells then crawl against the direction of CBF and probe BMECs for sites permissive for diapedesis. Diapedesis eventually takes place either paracellularly through bi-/tricellular endothelial junctions or transcellularly by inducing the formation of pores within ECs. WSS Wall shear stress, PGSL-1 P-selectin glycoprotein ligand-1, VLA-4 very-late antigen 4, VCAM-1 vascular cell adhesion molecule-1, LFA1 lymphocyte function-associated antigen 1, ICAM-1 Intercellular adhesion molecule 1, PVS perivascular space, WSS Wall shear stress, BM basement membrane

very-late antigen 4 (VLA-4) and lymphocyte function-associated antigen 1 (LFA-1) on leukocytes), and their respective ligands [151–153]. During inflammatory insults such as in MS, selectin and integrin expression increases in both BMECs and leukocytes [154]. In fact, inhibiting T cell adhesion by blocking the interaction of α_4 -integrins with brain endothelial vascular cell adhesion molecule-1 (VCAM-1) through natalizumab is currently one of the most effective treatments for relapsing–remitting MS [8, 155, 156]. The initial tethering is followed by rolling of the immune cell along the vascular wall with reduced speed. Interestingly, rolling does not seem to be required for successful T cell migration [58, 152, 157]. A direct capture mechanism of T cells, mediated by

α_4 -integrins/VCAM-1, has been shown predominantly under non-inflammatory conditions and might be specific for migration into the CNS [158–161].

During rolling, immune cells experience a tangential force caused by WSS, as well as a rotational torque due to their rolling motion, resulting in shear forces dragging on newly formed adhesion complexes [162]. Surprisingly, these forces were found to stabilize adhesive interactions rather than opposing them. This discovery led to the definition of ‘catch bonds’—a bond whose lifetime increases with rising tension, as opposed to ‘slip bonds’ whose interaction is destabilized by force [58, 163–165] (Fig. 2, insert). Catch bond formation is essential for the interaction of selectins and integrins whose lifetime is maximal

in a force range of 10–30 pN [57, 61, 62]. Application of lateral force in form of WSS is thought to accelerate the conformational transition of integrins into an extended high-affinity state and thus facilitate immune cell arrest in an outside-in manner [166–168].

Rolling at reduced velocity enables immune cells to probe the endothelial surface for chemokines [58]. These chemokines bind to chemokine receptors on the immune cell surface, delivering a G-protein coupled receptor (GPCR)-mediated inside-out signal to α - and β -integrins. GPCR signalling increases the affinity of integrins for their endothelial ligands via activation of the cytoskeletal adaptors talin and kindlin in the immune cell [169]. Interaction pairs include LFA-1 (T cell) and intercellular adhesion molecule-1/2 (ICAM-1/2) (ECs), as well as VLA-4 (T cell) and VCAM-1 (EC) [152, 153, 167, 170]. This is further facilitated by integrin clustering into focal adhesions and ultimately results in integrin-mediated immune cell arrest, polarization, and firm adhesion on the endothelium [58]. Catch bonds again play a crucial role in this process as integrin activation by chemokines alone is insufficient to stimulate adhesiveness [62, 171]. In summary, the adhesion of immune cells to the post-capillary bed is facilitated by both WSS and intercellular forces, together stabilizing the interaction and allowing for cell migration to occur.

T cell crawling on BMECs is directed against the flow

While adhering to the vascular wall, lymphocytes polarize with respect to the blood flow by adopting an elongated morphology with a wide and flat F-actin-rich lamellipodium at the leading edge and a tail-like projected uropod at the trailing edge [160, 172]. Crawling is an active movement facilitated by Rho GTPases [8, 58], involves the formation of additional adhesive contacts, and occurs at a highly reduced velocity of a few μm per minute [173, 174]. Forward propulsion along the endothelium is driven by actin polymerization and myosin II contractility (retrograde flow), which together push the lamellipodium forward, parallel to the surface [58, 63]. Thereby, the actin cytoskeleton generates internal forces of up to 0.6 nN [63], which are transferred to integrins via cytoskeletal adaptors such as talin, increasing the ability to form catch bonds [172]. This actin-talin-integrin linkage, however, is not absolute, leading to varying degrees of slippage [175]. This has been termed a ‘molecular clutch’ and is exploited by the cell to dynamically adjust migration speed and direction in response to changes in substrate adhesiveness and matrix stiffness [175, 176]. Enhanced stiffness, as found in inflammation and aging [148], has been linked to more persistent directional crawling [177] and increased efficiency of transmigration [178].

T cell crawling along the cerebral post-capillary venules is unique in two aspects. Firstly, crawling was observed in vivo to be directed preferentially against the direction of blood flow [8, 138, 152, 179]. This has been linked to the presence of shear forces and the adhesion molecules ICAM-1 and ICAM-2 [161, 172]. T cells can also crawl downstream by engagement of VLA-4 and VCAM-1, but prefer upstream migration under shear rates above 400 s^{-1} [172]. Interestingly, in the absence of VLA-4 or VCAM-1, T cells still adhere and crawl against the flow, but do not maintain directionality after the flow is terminated [161]. VLA-4 thus appears to be required for migratory memory, the capacity to remember directionality, under WSS [180].

Secondly, T cell crawling at the brain post-capillary venules takes place over significantly longer distances, frequently exceeding $150\text{ }\mu\text{m}$, which is likely due to the unique high barrier integrity rarely allowing for diapedesis even under inflammatory conditions [160]. In contrast to rapid rolling and tethering, crawling takes on average several minutes and rarely up to hours [138, 160, 161]. On primary mouse BMECs, 95% of T cells crawled or transmigrated within half an hour, whereas 5% arrested and remained immobile [160]. Similarly, intravital imaging of autoreactive T cells showed 80–85% of cells crawled and extravasated from the leptomeningeal vessels within 30 min [138]. The purpose and mechanism of preferential crawling against the flow at the BBB still remain elusive, and it has yet to be determined whether this phenomenon is immune cells specific and how frequently this event occurs in other tissues [179].

Diapedesis by the path of least resistance

Immune cell migration across BMECs occurs either paracellularly through endothelial junctions or transcellularly by inducing the formation of pore-like structures in the ECs themselves [152]. In search for permissive sites, leukocytes sense endothelial substrate stiffness, mainly defined by their F-actin density [148, 181]. Several factors determine which route will be preferred. High ICAM-1 levels, increased caveolin-1 expression, and chemokine presentation have all been linked to transcellular migration [182–186]. On the other hand, with high stress fibre density and lower levels of inflammation with intermediate ICAM-1 expression might favour paracellular diapedesis [182, 187]. Although increased stiffness appears to facilitate crawling, diapedesis of T cells locally takes place at the path of least mechanical resistance [182, 188]. Transcellular migration thus primarily occurs at the cell periphery, where F-actin density is highly heterogeneous [58, 182, 189]. As multiple forces and chemical signals act in concert, plus in vivo studies are sparse, it is challenging

to predict which migration route is preferred in neuro-inflammatory conditions like MS, AD, or during aging.

Finally, diapedesis takes place within seconds to minutes [190], during which both leukocytes and ECs undergo drastic cytoskeletal remodelling. Leukocytes adopt an elongated shape and drastically reduce stiffness by inducing actin disassembly and breaking down vimentin intermediate filaments and microtubules [189]. Once the leading edge of the leukocyte has breached the endothelial monolayer, the elongated leukocyte nucleus squeezes through the endothelium with lateral forces in the order of 60 nN to extrude through the narrow space [58]. This is sensed by the endothelial cytoskeleton, which responds with the disassembly of filaments to withstand these forces and allows for transmigration to succeed [189]. Leukocyte squeezing also triggers the formation of an F-actin ring surrounding the migrating immune cell. This ring limits pore size and prevents plasma leakage, allowing the endothelium to preserve its low permeability to macromolecules [189, 191]. Upon successful diapedesis, BMECs respond to the sudden loss of tension and utilize ventral lamellipodia to close the gap [189]. During transmigration, ECs and leukocytes exchange surface proteins [192, 193], which might draw additional immune cells to these sites, creating a hotspot for transmigration [194]. However, it is unclear whether the endothelium returns to its previous state eventually or retains long-lasting imprints at these sites.

The unique structural properties of the BBB would suggest that mechanical force transmission and response are distinct from those observed in other tissues, and might help explain molecular mechanisms directing transcellular or paracellular diapedesis. However, no studies have yet reported mechanical properties of diapedesis across BMECs.

Mechanosensory elements of endothelial cells

Considering the wide variety of mechanical stresses at the BBB, BMECs express a variety of mechanosensory complexes, which convert physical stress into biochemical signals (Fig. 3). Here we focus on cell–cell junction complexes and mechanosignalling through receptors and ion channels of BMECs, highlight their relevance in immune cell migration across the BBB, and changes during inflammation and aging. Importantly, we try to distinguish mechanosensing elements, which were shown to respond to mechanical forces directly, from mechanosensitive players, which may be activated indirectly by force transmission and biochemical signals downstream of primary mechanosensing components or actin fibres.

Cell–cell junctions

Cell–cell junctions play a crucial role in endothelial barrier function and have thus adapted multiple ways to respond to mechanical forces (Fig. 3) [4, 195]. Formation of stable AJs requires intracellular coupling of VE-cadherin to actomyosin through α -catenin [196, 197]. Both external forces on VE-cadherin and internal forces generated by actomyosin pull α -catenin into an extended conformation, leading to the recruitment of vinculin and other binding partners [4, 196, 197]. Vinculin further stabilizes the unfolded conformation of α -catenin, leading to enhanced cell–cell adhesion and F-actin polymerization [40, 195, 196]. Increasing force strengthens the binding (catch-bond) between the cytoskeleton and junctional partners vinculin, talin, and α -catenin upon formation of dimers/multimers (order of 10 pN), while e.g. a bond between monomeric α -catenin and F-actin disrupts under increasing tension (slip-bond) [198–201]. The actin cytoskeleton of ECs also directly responds to tension via polymerization and interaction with actin filament-associated proteins, leading to stress fibre formation, increased stiffness, junctional changes and activation of focal adhesions [40, 202].

Extracellular detachment forces between VE-cadherin proteins (Ca^{2+} -dependent) range from 15 to 150 pN depending on their cumulative interactions and on the applied stretch, which extends their lifetime (catch-bonds) [65, 203]. Cadherins are also thought to be mechanosensitive themselves, acting in concert with PECAM-1 and VEGF receptors to initiate further downstream signalling including PI3K/Akt and integrin/RhoA [6, 40, 64, 76, 204, 205]. Mechanosensing by PECAM-1 regulates force-dependent cellular stiffening [204] and PECAM-1 is critically involved in the regulation of BBB junctional integrity [206], WSS sensation [207], integrin activation, cytoskeletal rearrangement [195, 208] and TEM [209]. Loss of PECAM-1 increases vascular permeability of the BBB, but prevents paracellular transmigration, suggesting that intact (to a certain degree) cell junctions and PECAM-1 mechanosignalling are required for paracellular diapedesis [8, 185, 206]. On the other hand, in pathologies with disturbed WSS, such as atherosclerosis and ischemic stroke, PECAM-1 is increased and defined as a main contributor to atherosclerotic lesions, neutrophil invasion, and post-ischemic neuroinflammation [210, 211].

BM stiffness and WSS regulate the force on focal adhesions and TJs by traction force, hence mechanosensory properties have also been shown for TJ proteins. [6, 212]. Under physiological laminar WSS, mechanosensing via occludin promotes vascular integrity by recruiting further occludin and ZO-1, whereas disturbed flow causes loss of occludin and barrier




Fig. 3 Biomechanical signalling via mechanosensing/sensitive elements at the brain endothelium. Interacellular tension is sensed by junctional molecules such as PECAM-1, VE-cadherin, and catenins. Interactions between PECAM-1, VE-cadherin, Piezo1, S1PR1, and VEGFR2 regulate barrier stability. Integrin signalling transmits ECM stiffness and stretch to the actin cytoskeleton and ion channels. Transient receptor potential (TRP) ion channels may be activated by direct mechanical force transfer from primary mechanosensing components (integrins) or via intermediate proteins (CD98) or cytoskeletal adaptors. Alternatively, they may be activated via biochemical signalling downstream of primary mechanosensing elements, involving synthesis and modification of intracellular messengers (DAG and Ca²⁺). WSS wall shear stress, S1P/S1PR1 sphingosine-1-phosphate receptor 1, VEGFR2 vascular endothelial growth factor receptor 2, Rac1 Rac family small GTPase 1, Cdc42 cell division cycle 42, JAMs junctional adhesion molecules, PECAM-1 platelet and endothelial cell adhesion molecule 1, α/β cat catenins, ZO zonula occludens, PLC phospholipase C, DAG diacylglycerol, PIP2 Phosphatidylinositol 4,5-bisphosphate, Ca²⁺ calcium, ER endoplasmic reticulum

dysfunction [213]. In addition, the TJ adaptor protein ZO-1 also regulates tension acting on VE-cadherin in response to WSS, similar to the function of α-catenin at AJs [205, 212, 214]. Together these components enable the dynamic response of junctional proteins to mechanical forces, which are required for endothelial barrier formation and maintenance [196, 215], and cell orientation in the direction of flow [195]. In neurological pathologies, both mechanical forces as well as organization and expression of cell–cell junctions change drastically. However, it remains challenging to create a causative relationship between these two

components, as other factors, such as cytokine signalling, majorly affect junctional integrity [216].

Mechanosensitive receptors and ion channels

Both ECs and migrating immune cells display a wide range of receptors and ion channels responding to mechanical stress, although mechanisms of their localization and activation are often not fully understood [217]. Mechanosensitive proteins can be activated by direct stress applied to the lipid bilayer, opening e.g. the ion channel through membrane tension, or via additional linked components outside the membrane [218, 219].

Activated proteins often undergo large conformational changes, such as unfolding of domains in talin, which are reversed upon force dissipation [220, 221]. Of note, translation of mechanical signals into cellular alterations occurs on different time scales e.g. with integrin activation (seconds to minutes) and longer duration for cytoskeletal rearrangements (minutes to hours) [220].

Tyrosine kinase receptors on ECs, including vascular endothelial growth factor receptor 2 (VEGFR2) and Tie-2, become phosphorylated under mechanical stretch, which triggers ligand-independent signalling [222, 223]. Importantly, VEGFR2 has multiple phosphorylation sites, which can regulate vascular permeability through VE-cadherin phosphorylation, but also proliferation and migration [224]. As a result, mechanical cues can regulate angiogenesis via VEGFR2-signalling [225]. ECM stiffness regulates the angiogenic potential via yes-associated protein (YAP)-Dil4-Notch1-VEGFR signalling, and promotes pro-angiogenic factors on softer vs. stiffer substrates [226]. YAP/transcriptional coactivator with PDZ-binding motif (TAZ) activation also induces an inflammatory endothelial phenotype under disturbed blood flow, hence inhibition of this pathway has been effective against atherosclerotic lesion development [227, 228]. Exploration of these tightly connected mechanosensitive pathways for therapeutic approaches could reinstate functional angiogenesis in cerebral vessels with stiffer BM and support areas of disturbed or reduced WSS such as in elderly, AD and ischemia [229].

Mechanosensing GPCRs change conformation under mechanical stress [230]. For example, endothelial GPR68 is activated by high WSS in small diameter arteries, triggering flow-mediated dilation [231, 232]. Interestingly, GPR68 is also activated by acidosis, a lowered blood pH, which can occur through prolonged lack of oxygen and CO₂ accumulation [233]. However, CO₂-induced CBF increase was not changed in mice lacking GPR68, thus the role of GPR68 for H⁺ sensing at cerebral arteries could be less prominent [234, 235]. Sphingosine-1-phosphate receptor 1 (S1PR1) is another mechanosensitive GPCR expressed in BMECs but also widely throughout other tissues [236]. Vascular S1PR1 can be activated by WSS and circulating S1P resulting in vasodilation and maintenance of endothelial barrier stability [237]. Of note, S1PR modulators, such as fingolimod (S1PR1,3,4,5 modulator) and siponimod (S1PR1,5), are approved as disease-modifying treatment in MS, and tested in clinical trials for ischemic and hemorrhagic stroke [238, 239]. Fingolimod also reduces VCAM-1 expression and stabilizes claudin-5 levels on BMECs [240]. In line with S1PR1s mechanosensitive functions in vasodilation and vascular network maintenance, prolonged treatment with fingolimod resulted in hypertension in mice [241] and

patients with chronic inflammatory demyelinating polyradiculoneuropathy [242]. In conclusion, mechanosensitive GPCRs present a promising target group for multiple CNS diseases, however, their widespread expression and functional spectrum may limit therapeutic potential.

Mechanosensitive ion channels are primarily cation permeable and regulate ion flux to initiate Ca²⁺ signalling, alter the membrane potential, and regulate Mg²⁺ homeostasis [36, 40, 243]. Due to their rapid action and the fast propagation of Ca²⁺ waves, ion channel activity is usually among the first steps in mechanical signalling [40, 244]. This Ca²⁺ signalling is critical for maintaining the low permeability of the BBB [245], whereas dysregulation of intracellular Ca²⁺ due to high mechanical stress or pathological conditions can compromise BBB integrity [245–248]. For example, Ca²⁺ influx through the nonselective cation channel Piezo1 is involved in flow-induced cell alignment, vasodilation of cerebral capillaries and angiogenesis [40, 249–251]. Overstimulation of Piezo1 due to disturbed flow conditions in atherosclerosis and hypertension, however, causes the breakdown of AJs, actin remodelling and inflammation [78, 252–255]. The underlying mechanism is a direct connection between Piezo1 and VE-cadherin as well as PECAM-1 regulating junction integrity [256].

Mechanosensitive ion channels also include the diverse family of transient receptor potential (TRP) ion channels, which mediate large parts of the mechanical stress response experienced by BMECs. They fulfil a multitude of functions in immune cell migration and barrier function and are involved in both progression and recovery of several neurological and vascular-related pathologies [35, 36, 257]. The following section will highlight the functional diversity of this family of ion channels.

Mechanosensitive TRP channels

TRP channels are widely expressed throughout the CNS and are involved in cellular response to a variety of external cues, including mechanical signals [258] (Table 2). Based on sequence homology, they are grouped into six subfamilies termed canonical (TRPC1-7), vanilloid (TRPV1-6), ankyrin (TRPA1), polycystic (TRPP1-3), melastatin (TRPM1-8) and mucolipin (TRPML1-3) [36]. TRP channels can directly or indirectly trigger Ca²⁺ signalling [40, 245] and can also be Ca²⁺-regulated themselves via Ca²⁺-calmodulin or direct Ca²⁺ binding [40]. Although TRP channels are commonly involved in mechanosignalling cascades, likely none of them are mechanosensing themselves [259]. Mechanisms for direct mechanical activation, such as deformation due to membrane stretch, have been proposed, but lack sufficient evidence [260]. TRP channels are thus most likely activated by mechanical transduction via interaction with mechanosensing

Table 2 Mechanosensitive TRP channels of vascular ECs and leukocytes

Channel	Cell type	Upstream components	Reference(s)
TRPA1	BMECs, T cells, B cells, macrophages	[Ca ²⁺] _i	[262–264]
TRPC1	cBAECs, mBMECs, T cells, B cells, macrophages, neutrophils	PLC, DAG, [Ca ²⁺] _i	[246, 265–268]
TRPC3 ^a	cBAECs, mBMECs, T cells, B cells, macrophages, neutrophils	PLC, IP ₃	[246, 265, 266, 269–272]
TRPC5 ^a	cBAECs, mBMECs, macrophages	PLC, PIP ₂	[246, 266, 273, 274]
TRPC6	cBAECs, macrophages, neutrophils	DAG, Gq _α , PECAM-1	[266, 275–278]
TRPM4 ^a	BMECs, T cells, macrophages	IP ₃ , Ca ²⁺ -calmodulin	[279–282]
TRPM7 ^a	HUVECs, T cells, macrophages		[265, 283]
TRPP1	HUVECs		[284]
TRPP2	BMECs	glycocalyx, IP ₃	[247, 261]
TRPV1 ^a	BMECs, T cells, macrophages, neutrophils		[285–288]
TRPV2	BMECs, T cells, B cells, macrophages, neutrophils	phosphatidylinositol-3 kinase, Akt	[285, 286, 288–290]
TRPV4	BMECs, T cells, macrophages, neutrophils	β ₁ -integrins, CD98, Piezo1, PLC, PIP ₂ , Ca ²⁺ -calmodulin, arachidonic acid metabolites	[78, 285, 286, 291–295]
TRPV5 ^a	T cells, granulocytes		[296]
TRPV6 ^a	Jurkat T cells, granulocytes		[297]

Expression of mechanosensitive TRP channels in human primary vascular ECs (BMECs where data available) and leukocytes, and associated suggested upstream signalling components; BMECs–(human) brain microvascular endothelial cells; [Ca²⁺]_i–intracellular Ca²⁺; cBAECs–cultured brain artery endothelial cells; DAG–diacylglycerol; HUVECs–human umbilical cord endothelial cells; mBMECs–mouse brain microvessel endothelial cells; PIP₂–phosphatidylinositol-4,5-bisphosphate; PLC–phospholipase C

^a Mechanosensitive properties derived from cell types other than ECs or leukocytes

proteins, such as β₁-integrins and the glycocalyx [40, 261]. Alternatively, force transmission may activate TRP channels via intermediary membrane proteins and the cytoskeleton [40]. Table 2 provides an overview of known mechanosensitive TRP channels and associated upstream components. For several TRP subunits mechanosensitive properties have not been shown specifically for vascular ECs, but are likely based on findings in other cell types (indicated by superscript **a** in Table 2).

TRPV4 is a key mediator of mechanical signalling

TRPV4 is a non-selective cation channel triggering downstream signalling via influx of extracellular Ca²⁺ [298] and is involved in mechanosignalling of various cell types, including BMECs [286], T cells [285], neutrophils [294] and macrophages [295]. TRPV4 has been proposed to react to diverse mechanical forces such as WSS [78, 299–302], circumferential stretch [303, 304], stiffness [305, 306], osmotic swelling [298, 307], shrinkage and surface expansion [298].

Endothelial TRPV4 is an important regulator for flow-induced vasodilation [299, 302], cell volume [243, 293], cytoskeletal remodelling and barrier function [248, 308, 309]. TRPV4 activation can occur within milliseconds in focal adhesions of ECs by force transfer from mechanosensing β₁-integrins via CD98 to TRPV4 [292, 310]. TRPV4 activation by β₁-integrins might explain why high ECM stiffness, such as during inflammation

or aging, causes endothelial dysfunction in a pathway that involves TRPV4-mediated Ca²⁺ signalling [305]. TRPV4 is also involved in flow-induced vasodilation and cytoskeletal remodelling via Piezo1. Upon mechanical stimulation, Piezo1 causes a local increase in intracellular Ca²⁺ and activates TRPV4 via Ca²⁺-calmodulin binding to its C-terminal calmodulin binding site [78, 298, 311]. TRPV4 then triggers a more prolonged Ca²⁺ signal, which leads to Piezo1-mediated disassembly of AJs under disturbed blood flow [78]. TRPV4 inhibition in a mouse model of haemorrhagic stroke reduced BBB dysfunction, likely due to reduced Piezo1 signalling [309, 312]. These findings suggest a regulatory role of TRPV4 in vascular permeability, which contributes to BBB disruption under altered mechanical conditions [308, 309]. This role in BBB function and fast activation via β₁-integrins render TRPV4 a potential contributor to immune cell crawling and diapedesis [248]. It remains to be determined at what stage of the migration cascade TRPV4-mediated signalling occurs.

TRPC channels form mechanosensitive homomeric and heteromeric channels

Channels of the TRPC subfamily are involved in stretch and shear stress sensing of ECs and migrating immune cells [313, 314]. Although the expression of TRPCs in primary human BMECs has not explicitly been shown, their frequent expression in other human endothelial

tissue [315] would suggest that they are also common at the BBB. This is further supported by findings of TRPC expression in cultured brain artery ECs (cBAECs) [266] and mouse brain microvessel ECs (mBMECs) [246, 316]. All TRPC channel complexes in humans can be activated by PLC-mediated hydrolysis of PIP_2 [278, 317]. This can directly lead to TRPC channel opening by binding of diacylglycerol (DAG) or removal of PIP_2 -dependent inhibition [317–319]. Alternatively, IP_3 may travel through the cytoplasm, bind to its receptor and trigger the release of Ca^{2+} from intracellular stores, which in turn activates TRPC channels by Ca^{2+} or Ca^{2+} -calmodulin binding [35, 260, 317, 320]. Activation by mechanical stress might thus involve mechanosensing GPCRs activating $G_{q/11}$ proteins, which in turn stimulate PLC and lead to TRPC channel opening further downstream [313].

Whereas TRPC1 is essential for cell migration and polarization [203, 243, 313], TRPC6 plays an important regulatory role in barrier-forming tissue [40, 209]. Endothelial TRPC6 channels mediate the intracellular Ca^{2+} signalling to allow leukocyte diapedesis [189, 209]. Weber et al. [209] showed that TRPC6-deficient ECs prevent paracellular diapedesis, whereas selective activation of TRPC6 by a DAG analogue allows for transmigration even when PECAM-1 is blocked. These results suggest that TRPC6 is activated downstream of PECAM-1-mediated mechanosensing and is required for the cytoskeletal adaptations in ECs enabling leukocytes passing through. The same study also showed that TRPC6 channels accumulate around PECAM-1 and promote translocation of intracellular stores of PECAM-1 and other adhesion molecules to the site of diapedesis [209]. TRPC6-induced Ca^{2+} signalling is likely also responsible for the formation of the F-actin ring surrounding migrating immune cells [189]. Under pro-inflammatory and disturbed flow conditions, TRPC6 expression and activation are elevated independently of diapedesis, leading to EC contraction and increased vascular permeability [321, 322].

TRPC channels function both as homomeric complexes and in heteromeric combinations, which differ in terms of activation characteristics, sensitivity and cation selectivity [40, 314, 323]. For example, vasorelaxation by ECs in response to high WSS is in part realized by heteromeric complexes of TRPC1 and TRPV4 [40, 314, 324]. The formation of heteromers might explain the multiple and often overlapping functionalities of TRP subunits [40, 314] and may be a major reason why most TRP knockouts in animal models do not show a detrimental phenotype [309]. How the formation of these complexes is regulated and how their functionalities differ from their homomeric counterparts requires further investigation.

TRPP, TRPM and TRPA1 channels regulate vascular tone, permeability and function

TRPP1 and TRPP2, also known as Polycystin-1 and -2, are nonselective cation channels involved in flow and pressure sensing of vascular ECs [323, 325]. Both subunits accumulate at primary cilia where they regulate vascular tone in response to WSS by inducing nitric oxide production [284, 325, 326]. Upon mechanical injury of BMECs, TRPP2 is involved in stress fibre formation and cytoskeletal remodelling leading to BBB dysfunction [245, 247]. Interestingly, TRPP2 may only be mechanosensitive in combination with other TRP subunits, including TRPP1, TRPC1 and TRPV4, once again highlighting the relevance of heteromeric TRP complexes [323, 325]. Accordingly, pressure sensing by ECs was shown to be dependent on relative expression of TRPP1 and TRPP2 [325], and heteromeric complexes of TRPV4 and TRPP2 are much more sensitive to laminar WSS than homomeric TRPV4 channels [327]. Changing composition of heteromeric TRP channels might thus present a useful way by which cells regulate their sensitivity to mechanical stress.

TRPM channels are widely expressed in the brain and are involved in many neurological disorders [35]. TRPM4 was shown to be activated by membrane stretch in vascular smooth muscle cells [280], but its role in BMECs still remains elusive. In vascular ECs, TRPM4 associates with Sulfonylurea receptor 1, which is upregulated in BMECs after ischemic stroke [35]. This causes continuous Na^+ influx and can lead to cell death, disintegration of capillaries and secondary bleeding [35, 328]. TRPM7 is another potentially mechanosensitive channel, which is activated by shear stress in fibroblasts [329], vascular smooth muscle cells [330] and mesenchymal stem cells [331]. Knockdown of TRPM7 in human umbilical cord ECs (HUVECs) prevents adhesion of ECs and promotes their growth and proliferation, suggesting a regulatory role in cell differentiation [283].

TRPA1 is involved in vasodilation of BMECs [262, 332] and is also expressed in T cells, where it promotes immune functions in response to mechanical stress and is implicated with several inflammatory conditions [263, 264]. TRPA1 can be activated by Ca^{2+} influx [35] and may even be mechanosensing itself. Although a recent study showed that TRPA1 in reconstituted proteoliposomes is insensitive to membrane stretch [259], another study revealed that single-channel currents of TRPA1 in artificial lipid bilayers respond to mechanical stretch in a pressure-dependent way [333]. This discrepancy might be explained by its redox state, as mechanosensing of TRPA1 was abolished under reducing conditions [333]. Clearly, TRP channel activation and function are not yet fully explained. Further studies should focus on the

effects of redox and other environmental conditions on TRP subunits, as these might help to better explain their activation, function and changes under pathological conditions.

Discussion and future perspectives

Mechanical stressors are crucial regulators of BBB function. Physiological WSS and BM stiffness support the formation of AJs and TJs and suppress inflammatory signalling, whereas alterations under pathological conditions lead to enhanced vascular permeability, endothelial dysfunction, reduced vascular reactivity and facilitate leukocyte infiltration. In this review, we took a holistic view of the various aspects of mechanical forces and mechanosensing at BMECs in different neuropathologies, which open new future intervention avenues.

Increasing evidence suggests that mechanisms of immune cell infiltration are not conserved across the body and depend on the tissue and immune cell type involved. Such cell-type-specific events offer great opportunities for therapeutic interventions. In MS, for example, Th1, Th17 and CD8⁺ T cells are thought to promote inflammation in the CNS, whereas Th2 cells are associated with a protective effect [334, 335]. Th2 cells show lower ability to form tethers and slings and thus egress to a much lesser extent under high WSS [336, 337]. T cell subtype-dependent behaviour has also been observed for crawling [338] and diapedesis [185], but the underlying molecular mechanisms governing these differences remain mostly undescribed [189]. Closer examination of the molecular players driving cell-type-specific events throughout migration could reveal novel therapeutic targets for the treatment of MS and other neuroinflammatory or age-related conditions.

TEM is controlled by a range of mechanosensory elements on both migrating leukocytes and the endothelium. It is not always clear whether certain mechanosensory proteins adapt to pathological conditions to maintain vascular integrity, or whether these conditions throw mechanosignalling events off-balance and thereby contribute to pathological hallmarks. For example, PECAM-1 mediates vasodilation to alleviate high shear stress and might be involved in restoring the BBB after disruption in MS [206, 339], but also contributes to immune cell infiltration and inflammation in atherosclerosis, stroke and MS [211, 339, 340]. Moreover, disturbed flow and inflammation frequently affect expression and localization of endothelial proteins, further altering their response [341]. Decreased surface expression of VE-cadherin during inflammation increases permeability and reduces the ability of ECs to respond to WSS [342]. To arrive at a better understanding of pathological effects on the mechanosensory response, future research will

require a more detailed exploration of mechanosensing and -sensitive proteins during neuroinflammation and aging.

Mechanosensitive TRP channels fulfil a wide range of functions and thus offer a plethora of potential drug interventions. Tremendous effort has been put into the development of specific TRP channel agonists and antagonists as drug targets for a range of diseases, including MS, Alzheimer's disease and stroke [343]. A recent study from our lab explores specifically the potential of TRPV4 inhibition to reduce leukocyte migration across the BBB in MS, however without integrating WSS or BM stiffness [248]. Although numerous potential compounds have entered clinical trials, several of them were eventually withdrawn due to lacking effectiveness and a variety of adverse effects [343, 344]. To effectively use TRP channels for drug discovery, a better understanding of channel expression and activity during pathology is required.

In conclusion, biomechanical signalling is an essential component of BBB function, yet many questions remain open on how these signals change during neuropathology. Continuing research in this field will provide new insights into the cumulative effects of mechanical alterations in WSS, BM stiffness and cell–cell interactions during immune cell migration. Ideally, such understanding would lead to the development of novel drug candidates selectively altering mechanosensory properties of immune cell subsets while leaving other functionalities unaffected, thus counteracting the problem of multifunctionality.

Abbreviations

Aβ	Amyloid beta
ACA	Anterior cerebral artery
AD	Alzheimer's disease
AJ	Adherens junction
BAEC	Brain artery endothelial cells
BBB	Blood–brain barrier
BM	Basement membrane
BMEC	Brain microvascular endothelial cell
CAA	Cerebral amyloid angiopathy
CBF	Cerebral blood flow
CCA	Common carotid artery
Cdc42	Cell division cycle 42
CNS	Central nervous system
CSF	Cerebrospinal fluid
DAG	Diacylglycerol
EC	Endothelial cell
ECM	Extracellular matrix
WSS	Wall shear stress
GPCR	G-protein coupled receptor
HSPGs	Heparan sulfate proteoglycans
HUVEC	Human umbilical cord endothelial cells
ICA	Internal carotid artery
ICAM-1	Intercellular adhesion molecule 1
IJV	Internal jugular vein
IP ₃	Inositol 1,4,5-trisphosphate
JAM	Junctional adhesion molecules
LBRC	Lateral border recycling compartment
LFA1	Lymphocyte function-associated antigen 1
mBMEC	Mouse brain microvessel endothelial cells

MRI	Magnetic resonance imaging
MS	Multiple sclerosis
NFκB	Nuclear factor kappa B subunit
PECAM-1	Platelet and endothelial cell adhesion molecule 1
PIP ₂	Phosphatidylinositol-4,5-bisphosphate
PLC	Phospholipase C
Rac1	Rac family small GTPase 1
ROS	Reactive oxygen species
S1P/S1PR1	Sphingosine-1-phosphate receptor 1
TAZ	Transcriptional coactivator with PDZ-binding motif
TEM	Transendothelial migration
TJ	Tight junction
TRP	Transient receptor potential
VCAM-1	Vascular cell adhesion molecule-1
VEGFR2	Vascular endothelial growth factor receptor 2
VLA-4	Very-late antigen 4
YAP	Yes-associated protein
ZO	Zonula occludens

Author contributions

CEH and DH designed and wrote the manuscript. AK created the figures and revised the manuscript. HEDV contributed to the design of the study, obtained research grants, and revised the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding

This work was funded by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant (ENTRAIN) (agreement No. 813294) to EdV.

Availability of data and materials

No datasets were generated or analysed during the current study.

Declarations

Ethics approval and consent to participate

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 7 October 2024 Accepted: 7 December 2024

Published online: 18 December 2024

References

- Obermeier B, Verma A, Ransohoff RM. The blood–brain barrier. In: Pittock SJ, Vincent A, editors. *Handbook of clinical neurology*. Amsterdam: Elsevier; 2016. p. 39–59.
- Varatharaj A, Galea I. The blood–brain barrier in systemic inflammation. *Brain Behav Immun*. 2017;60:1–12.
- Zlokovic BV. The blood–brain barrier in health and chronic neurodegenerative disorders. *Neuron*. 2008;57(2):178–201.
- Angulo-Urarte A, Van Der Wal T, Huveneers S. Cell–cell junctions as sensors and transducers of mechanical forces. *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 2020;1862(9):183316.
- Cucullo L, Hossain M, Puvanna V, Marchi N, Janigro D. The role of shear stress in blood–brain barrier endothelial physiology. *BMC Neurosci*. 2011;12:40.
- Conway DE, Breckenridge MT, Hinde E, Gratton E, Chen CS, Schwartz MA. Fluid shear stress on endothelial cells modulates mechanical tension across VE-cadherin and PECAM-1. *Curr Biol*. 2013;23(11):1024–30.
- Profaci CP, Munji RN, Pulido RS, Daneman R. The blood–brain barrier in health and disease: Important unanswered questions. *J Exp Med*. 2020. <https://doi.org/10.1084/jem.20190062>.
- Marchetti L, Engelhardt B. Immune cell trafficking across the blood–brain barrier in the absence and presence of neuroinflammation. *Vasc Biol*. 2020;2(1):H1–h18.
- Castro Dias M, Coisne C, Lazarevic I, Baden P, Hata M, Iwamoto N, et al. Claudin-3-deficient C57BL/6J mice display intact brain barriers. 2019. *Sci Rep*. <https://doi.org/10.1038/s41598-018-36731-3>.
- Iwamoto N, Higashi T, Furuse M. Localization of angulin-1/LSR and tricellulin at tricellular contacts of brain and retinal endothelial cells in vivo. *Cell Struct Funct*. 2014;39(1):1–8.
- Arts JGG, Mahlandt EK, Schimmel L, Grönloh MLB, van der Niet S, Klein B, et al. Endothelial focal adhesions are functional obstacles for leukocytes during basolateral crawling. *Front Immunol*. 2021;12:667213.
- Minagar A, Alexander JS. Blood–brain barrier disruption in multiple sclerosis. *Mult Scler*. 2003;9(6):540–9.
- Alexander JS, Zivadinov R, Maghzi A-H, Ganta VC, Harris MK, Minagar A. Multiple sclerosis and cerebral endothelial dysfunction: mechanisms. *Pathophysiology*. 2011;18(1):3–12.
- Pradeepkiran JA, Baig J, Islam MA, Kshirsagar S, Reddy PH. Amyloid-β and phosphorylated tau are the key biomarkers and predictors of Alzheimer's disease. *Aging Dis*. 2024. <https://doi.org/10.1433/AD.2024.0286>.
- Mehta RI, Mehta RI. The vascular-immune hypothesis of Alzheimer's disease. *Biomedicines*. 2023. <https://doi.org/10.3390/biomedicines11020408>.
- Greenberg SM, Bacskai BJ, Hernandez-Guillamon M, Pruzin J, Sperling R, van Veluw SJ. Cerebral amyloid angiopathy and Alzheimer disease— one peptide, two pathways. *Nat Rev Neurol*. 2020;16(1):30–42.
- Ho VS, Cenzer IS, Nguyen BT, Lee SJ. Time to benefit for stroke reduction after blood pressure treatment in older adults: a meta-analysis. *J Am Geriatr Soc*. 2022;70(5):1558–68.
- Cheng X, Potenza DM, Brenna A, Ajalbert G, Yang Z, Ming XF. Aging increases hypoxia-induced endothelial permeability and blood–brain barrier dysfunction by upregulating arginase-II. *Aging Dis*. 2024. <https://doi.org/10.1433/AD.2023.1225>.
- Binnewijzend MA, Benedictus MR, Kuijter JP, van der Flier WM, Teunissen CE, Prins ND, et al. Cerebral perfusion in the prodromal stages of Alzheimer's disease. *Eur Radiol*. 2016;26(2):506–14.
- Hays CC, Zlatar ZZ, Wierenga CE. The utility of cerebral blood flow as a biomarker of preclinical Alzheimer's disease. *Cell Mol Neurobiol*. 2016;36(2):167–79.
- Shimizu H, Ghazizadeh M, Sato S, Oguro T, Kawanami O. Interaction between beta-amyloid protein and heparan sulfate proteoglycans from the cerebral capillary basement membrane in Alzheimer's disease. *J Clin Neurosci*. 2009;16(2):277–82.
- Howe MD, McCullough LD, Urayama A. The role of basement membranes in cerebral amyloid angiopathy. *Front Physiol*. 2020;11:601320.
- Chabriat H, Joutel A, Dichgans M, Tournier-Lasserre E, Bousser MG, Cadasil. *Lancet Neurol*. 2009;8(7):643–53.
- Joutel A, Monet-Leprêtre M, Gosele C, Baron-Menguy C, Hammes A, Schmidt S, et al. Cerebrovascular dysfunction and microcirculation rarefaction precede white matter lesions in a mouse genetic model of cerebral ischemic small vessel disease. *J Clin Invest*. 2010;120(2):433–45.
- Joutel A, Haddad I, Ratelade J, Nelson MT. Perturbations of the cerebrovascular matrisome: a convergent mechanism in small vessel disease of the brain? *J Cereb Blood Flow Metab*. 2016;36(1):143–57.
- Saji N, Toba K, Sakurai T. Cerebral small vessel disease and arterial stiffness: tsunami effect in the brain? *Pulse*. 2016;3(3–4):182–9.
- Weller RO, Subash M, Preston SD, Mazanti I, Carare RO. Perivascular drainage of amyloid-beta peptides from the brain and its failure in cerebral amyloid angiopathy and Alzheimer's disease. *Brain Pathol*. 2008;18(2):253–66.
- Govindpani K, McNamara LG, Smith NR, Vinnakota C, Waldvogel HJ, Faull RL, Kwakowsky A. Vascular dysfunction in Alzheimer's disease: a prelude to the pathological process or a consequence of it? *J Clin Med*. 2019. <https://doi.org/10.3390/jcm8050651>.
- Katritsis D, Kaiktsis L, Chaniotis A, Pantos J, Efstathiopoulos EP, Marmarelis V. Wall shear stress: theoretical considerations and methods of measurement. *Prog Cardiovasc Dis*. 2007;49(5):307–29.
- Zabrodskaya Y, Paramonova N, Litovchenko A, Bazhanova E, Gerasimov A, Sitovskaya D, et al. Neuroinflammatory dysfunction of the blood–brain barrier and basement membrane dysplasia play a role in the

- development of drug-resistant epilepsy. *Int J Mol Sci.* 2023. <https://doi.org/10.3390/ijms241612689>.
31. Greiner T, Kipp M. What guides peripheral immune cells into the central nervous system? *Cells.* 2021. <https://doi.org/10.3390/cells10082041>.
 32. Hu D, Weiner HL. Unraveling the dual nature of brain CD8+ T cells in Alzheimer's disease. *Mol Neurodegener.* 2024;19(1):16.
 33. Smolders J, Heutink KM, Fransen NL, Remmerswaal EBM, Hombrink P, Ten Berge IJM, et al. Tissue-resident memory T cells populate the human brain. *Nat Commun.* 2018;9(1):4593.
 34. Ortiz GG, Pacheco-Moisés FP, Macías-Islas M, Flores-Alvarado LJ, Mireles-Ramírez MA, González-Renovato ED, et al. Role of the blood-brain barrier in multiple sclerosis. *Arch Med Res.* 2014;45(8):687–97.
 35. Huang Q, Wang X, Lin X, Zhang J, You X, Shao A. The role of transient receptor potential channels in blood-brain barrier dysfunction after ischemic stroke. *Biomed Pharmacother.* 2020;131:110647.
 36. Earley S, Brayden JE. Transient receptor potential channels in the vasculature. *Physiol Rev.* 2015;95(2):645–90.
 37. Vanlandewijck M, He L, Mäe MA, Andrae J, Ando K, Del Gaudio F, et al. A molecular atlas of cell types and zonation in the brain vasculature. *Nature.* 2018;554(7693):475–80.
 38. Wang X, Xu B, Xiang M, Yang X, Liu Y, Liu X, Shen Y. Advances on fluid shear stress regulating blood-brain barrier. *Microvasc Res.* 2020;128:103930.
 39. Gray KM, Stroka KM. Vascular endothelial cell mechanosensing: new insights gained from biomimetic microfluidic models. *Semin Cell Dev Biol.* 2017;71:106–17.
 40. Fels B, Kusche-Vihrog K. It takes more than two to tango: mechanosignaling of the endothelial surface. *Pflügers Arch Eur J Physiol.* 2020;472(4):419–33.
 41. Zhang X, Kim T-H, Thauland TJ, Li H, Majedi FS, Ly C, et al. Unraveling the mechanobiology of immune cells. *Curr Opin Biotechnol.* 2020;66:236–45.
 42. Samady H, Eshtehardi P, McDaniel MC, Suo J, Dhawan SS, Maynard C, et al. Coronary artery wall shear stress is associated with progression and transformation of atherosclerotic plaque and arterial remodeling in patients with coronary artery disease. *Circulation.* 2011;124(7):779–88.
 43. Hademenos GJ, Massoud TF. Biophysical mechanisms of stroke. *Stroke.* 1997;28(10):2067–77.
 44. Kamimura T, Aoki S, Nezu T, Eto F, Shiga Y, Nakamori M, et al. Association between carotid wall shear stress-based vascular vector flow mapping and cerebral small vessel disease. *J Atheroscler Thromb.* 2023;30(9):1165–75.
 45. Rastegari K, Mokhtari-Dizaji M, Harirchian MH, Hashemi H, Ayooobi Yazdi N, Saberi H. Biomechanical changes of the common carotid artery and internal jugular vein in patients with multiple sclerosis. *Ultrasonography.* 2023;42(1):100–10.
 46. Kamiya A, Bukhari R, Togawa T. Adaptive regulation of wall shear stress optimizing vascular tree function. *Bull Math Biol.* 1984;46(1):127–37.
 47. Wong AD, Ye M, Levy AF, Rothstein JD, Bergles DE, Searson PC. The blood-brain barrier: an engineering perspective. *Front Neuroeng.* 2013;6:7.
 48. Koutsouris AG, Tachmitzi SV, Batis N, Kotoula MG, Karabatsas CH, Tsironi E, Chatzoulis DZ. Volume flow and wall shear stress quantification in the human conjunctival capillaries and post-capillary venules in vivo. *Biorheology.* 2007;44(5–6):375–86.
 49. Ballermann BJ, Dardik A, Eng E, Liu A. Shear stress and the endothelium. *Kidney Int.* 1998;54:S100–8.
 50. Abaci HE, Shen Y-I, Tan S, Gerecht S. Recapitulating physiological and pathological shear stress and oxygen to model vasculature in health and disease. *Sci Rep.* 2014;4(1):4951.
 51. Cheng H, Chen X, Zhong J, Li J, Qiu P, Wang K. Label-free measurement of wall shear stress in the brain venule and arteriole using dual-wave-length third-harmonic-generation line-scanning imaging. *Opt Lett.* 2022;47(21):5618–21.
 52. Grifno GN, Farrell AM, Linville RM, Arevalo D, Kim JH, Gu L, Searson PC. Tissue-engineered blood-brain barrier models via directed differentiation of human induced pluripotent stem cells. *Sci Rep.* 2019;9(1):13957.
 53. Candiello J, Cole GJ, Halfter W. Age-dependent changes in the structure, composition and biophysical properties of a human basement membrane. *Matrix Biol.* 2010;29(5):402–10.
 54. Gordon E, Schimmel L, Frye M. The importance of mechanical forces for in vitro endothelial cell biology. *Front Physiol.* 2020;11:684.
 55. Kohn JC, Lampi MC, Reinhart-King CA. Age-related vascular stiffening: causes and consequences. *Front Genet.* 2015. <https://doi.org/10.3389/fgene.2015.00112/full>.
 56. Xue C, Zhang T, Xie X, Zhang Q, Zhang S, Zhu B, et al. Substrate stiffness regulates arterial-venous differentiation of endothelial progenitor cells via the Ras/Mek pathway. *Biochim Biophys Acta Mol Cell Res.* 2017;1864(10):1799–808.
 57. Marshall BT, Long M, Piper JW, Yago T, McEver RP, Zhu C. Direct observation of catch bonds involving cell-adhesion molecules. *Nature.* 2003;423(6936):190–3.
 58. Huse M. Mechanical forces in the immune system. *Nat Rev Immunol.* 2017;17(11):679–90.
 59. Zhu C, McEver RP. Catch bonds: physical models and biological functions. *Mol Cell Biomech.* 2005;2(3):91.
 60. Morikis VA, Chase S, Wun T, Chaikof EL, Magnani JL, Simon SI. Selectin catch-bonds mechanotransduce integrin activation and neutrophil arrest on inflamed endothelium under shear flow. *Blood.* 2017;130(19):2101–10.
 61. Kong F, García AJ, Mould AP, Humphries MJ, Zhu C. Demonstration of catch bonds between an integrin and its ligand. *J Cell Biol.* 2009;185(7):1275–84.
 62. Chen W, Lou J, Zhu C. Forcing switch from short- to intermediate- and long-lived states of the aa domain generates LFA-1/ICAM-1 catch bonds. *J Biol Chem.* 2010;285(46):35967–78.
 63. Valignat MP, Theodoly O, Gucciardi A, Hogg N, Lellouch AC. T lymphocytes orient against the direction of fluid flow during LFA-1-mediated migration. *Biophys J.* 2013;104(2):322–31.
 64. Baumgartner W, Hinterdorfer P, Ness W, Raab A, Vestweber D, Schindler H, Drenkhahn D. Cadherin interaction probed by atomic force microscopy. *Proc Natl Acad Sci USA.* 2000;97(8):4005–10.
 65. Arslan FN, Eckert J, Schmidt T, Heisenberg CP. Holding it together: when cadherin meets cadherin. *Biophys J.* 2021;120(19):4182–92.
 66. Roux E, Bougaran P, Dufourcq P, Couffignal T. Fluid shear stress sensing by the endothelial layer. *Front Physiol.* 2020. <https://doi.org/10.3389/fphys.2020.00861>.
 67. Dewey CF Jr, Bussolari SR, Gimbrone MA Jr, Davies PF. The dynamic response of vascular endothelial cells to fluid shear stress. *J Biomech Eng.* 1981;103(3):177–85.
 68. Rochford KD, Cummins PM. In vitro cell models of the human blood-brain barrier: demonstrating the beneficial influence of shear stress on brain microvascular endothelial cell phenotype. In: Barichello T, editor. *Blood-brain barrier*. New York: Springer; 2019. p. 71–98.
 69. Krizanac-Bengez L, Mayberg MR, Janigro D. The cerebral vasculature as a therapeutic target for neurological disorders and the role of shear stress in vascular homeostasis and pathophysiology. *Neurol Res.* 2004;26(8):846–53.
 70. Liu Z, Ling SD, Liang K, Chen Y, Niu Y, Sun L, et al. Viscoelasticity of ECM and cells—origin, measurement and correlation. *Mechanobiol Med.* 2024;2(4):100082.
 71. Dessalles CA, Leclech C, Castagnino A, Barakat AI. Integration of substrate- and flow-derived stresses in endothelial cell mechanobiology. *Commun Biol.* 2021;4(1):764.
 72. Tran M, Heo C, Lee LP, Cho H. Human mini-blood-brain barrier models for biomedical neuroscience research: a review. *Biomater Res.* 2022;26(1):82.
 73. Kim J, Shin S-A, Lee CS, Chung HJ. An improved in vitro blood-brain barrier model for the evaluation of drug permeability using transwell with shear stress. *Pharmaceutics.* 2024;16(1):48.
 74. Molins Gutiérrez G, Martorell J, Salazar-Martin AG, Balcells M. A dynamic, in vitro BBB model to study the effects of varying levels of shear stress. In: Stone N, editor. *The blood-brain barrier: methods and protocols*. New York: Springer; 2022. p. 175–90.
 75. Korte N, Nortley R, Attwell D. Cerebral blood flow decrease as an early pathological mechanism in Alzheimer's disease. *Acta Neuropathol.* 2020;140(6):793–810.
 76. Baeyens N, Nicoli S, Schwartz MA. Vascular remodeling is governed by a VEGFR3-dependent fluid shear stress set point. *Elife.* 2015;4:e04645.
 77. Garcia-Polite F, Martorell J, Del Rey-Puech P, Melgar-Lesmes P, O'Brien CC, Roquer J, et al. Pulsatility and high shear stress deteriorate barrier

- phenotype in brain microvascular endothelium. *J Cereb Blood Flow Metab.* 2017;37(7):2614–25.
78. Swain SM, Liddle RA. Piezo1 acts upstream of TRPV4 to induce pathological changes in endothelial cells due to shear stress. *J Biol Chem.* 2021;296:100171.
 79. Pasi M, Cordonnier C. Clinical relevance of cerebral small vessel diseases. *Stroke.* 2020;51(1):47–53.
 80. Pasi M, Charidimou A, Boulouis G, Auril E, Ayres A, Schwab KM, et al. Mixed-location cerebral hemorrhage/microbleeds: Underlying microangiopathy and recurrence risk. *Neurology.* 2018;90(2):e119–26.
 81. Goeldlin MB, Mueller M, Siepen BM, Zhang W, Ozkan H, Locatelli M, et al. CADMUS. *Neurology.* 2024;102(1):e207977.
 82. Joutel A, Faraci FM. Cerebral small vessel disease. *Stroke.* 2014;45(4):1215–21.
 83. Wartolowska KA, Webb AJS. Midlife blood pressure is associated with the severity of white matter hyperintensities: analysis of the UK biobank cohort study. *Eur Heart J.* 2021;42(7):750–7.
 84. Bartstra JW, van den Beukel T, Kranenburg G, Geurts LJ, den Harder AM, Witkamp T, et al. Increased intracranial arterial pulsatility and microvascular brain damage in pseudoxanthoma elasticum. *Am J Neuroradiol.* 2024;45(4):386–92.
 85. Lu W, Yu C, Wang L, Wang F, Qiu J. Perfusion heterogeneity of cerebral small vessel disease revealed via arterial spin labeling MRI and machine learning. *Neuroimage Clin.* 2022;36:103165.
 86. Jiang X, Andjelkovic AV, Zhu L, Yang T, Bennett MVL, Chen J, et al. Blood-brain barrier dysfunction and recovery after ischemic stroke. *Prog Neurobiol.* 2018;163–164:144–71.
 87. Chang E, O'Donnell ME, Barakat AI. Shear stress and 17beta-estradiol modulate cerebral microvascular endothelial Na-K-Cl cotransporter and Na/H exchanger protein levels. *Am J Physiol Cell Physiol.* 2008;294(1):C363–71.
 88. Zhang M, Liu Q, Meng H, Duan H, Liu X, Wu J, et al. Ischemia-reperfusion injury: molecular mechanisms and therapeutic targets. *Signal Transduct Target Ther.* 2024;9(1):12.
 89. Ashby JW, Mack JJ. Endothelial control of cerebral blood flow. *Am J Pathol.* 2021;191(11):1906–16.
 90. Chen D, Li L, Wang Y, Xu R, Peng S, Zhou L, Deng Z. Ischemia-reperfusion injury of brain induces endothelial-mesenchymal transition and vascular fibrosis via activating let-7i/TGF- β 1 double-negative feedback loop. *FASEB J.* 2020;34(5):7178–91.
 91. Sun D, Ma J, Du L, Liu Q, Yue H, Peng C, et al. Fluid shear stress induced-endothelial phenotypic transition contributes to cerebral ischemia-reperfusion injury and repair. *APL Bioeng.* 2024. <https://doi.org/10.1063/5.0174825>.
 92. Protzmann J, Jung F, Jakobsson L, Fredriksson L. Analysis of ischemic stroke-mediated effects on blood–brain barrier properties along the arteriovenous axis assessed by intravital two-photon imaging. *Fluids Barriers CNS.* 2024;21(1):35.
 93. Latour LL, Kang DW, Ezzeddine MA, Chalela JA, Warach S. Early blood-brain barrier disruption in human focal brain ischemia. *Ann Neurol.* 2004;56(4):468–77.
 94. Schuff N, Matsumoto S, Kmiecik J, Studholme C, Du A, Ezekiel F, et al. Cerebral blood flow in ischemic vascular dementia and Alzheimer's disease, measured by arterial spin-labeling magnetic resonance imaging. *Alzheimers Dement.* 2009;5(6):454–62.
 95. Tomoto T, Tarumi T, Zhang R. Central arterial stiffness, brain white matter hyperintensity and total brain volume across the adult lifespan. *J Hypertens.* 2023. <https://doi.org/10.1097/HJH.0000000000003404>.
 96. Tarumi T, Zhang R. Cerebral blood flow in normal aging adults: cardiovascular determinants, clinical implications, and aerobic fitness. *J Neurochem.* 2018;144(5):595–608.
 97. Law M, Saindane AM, Ge Y, Babb JS, Johnson G, Mannon LJ, et al. Microvascular abnormality in relapsing-remitting multiple sclerosis: perfusion mr imaging findings in normal-appearing white matter. *Radiology.* 2004;231(3):645–52.
 98. Adhya S, Johnson G, Herbert J, Jaggi H, Babb JS, Grossman RI, Inglese M. Pattern of hemodynamic impairment in multiple sclerosis: dynamic susceptibility contrast perfusion MR imaging at 3.0 T. *Neuroimage.* 2006;33(4):1029–35.
 99. Senzaki K, Okada Y, Ochi H, Ochi M, Takei SI, Miura S, et al. Vascular endothelial dysfunction associated with severity in multiple sclerosis. *Multiple Scler Related Disorders.* 2021. <https://doi.org/10.1016/j.msard.2021.103135>.
 100. Mokhber N, Shariatzadeh A, Avan A, Saber H, Babaei GS, Chaimowitz G, Azarpazhooh MR. Cerebral blood flow changes during aging process and in cognitive disorders: a review. *Neuroradiol J.* 2021;34(4):300–7.
 101. Sankar SB, Pybus AF, Liew A, Sanders B, Shah KJ, Wood LB, Buckley EM. Low cerebral blood flow is a non-invasive biomarker of neuroinflammation after repetitive mild traumatic brain injury. *Neurobiol Dis.* 2019;124:544–54.
 102. Chen Y, Shen F, Liu J, Yang GY. Arterial stiffness and stroke: de-stiffening strategy, a therapeutic target for stroke. *Stroke Vasc Neurol.* 2017;2(2):65–72.
 103. Song J, Zhang X, Buscher K, Wang Y, Wang H, Di Russo J, et al. Endothelial basement membrane laminin 511 contributes to endothelial junctional tightness and thereby inhibits leukocyte transmigration. *Cell Rep.* 2017;18(5):1256–69.
 104. Thomsen MS, Routh LJ, Moos T. The vascular basement membrane in the healthy and pathological brain. *J Cereb Blood Flow Metab.* 2017;37(10):3300–17.
 105. Yao Y. Basement membrane and stroke. *J Cereb Blood Flow Metab.* 2019;39(1):3–19.
 106. Kretschmer M, Rüdiger D, Zahler S. Mechanical aspects of angiogenesis. *Cancers.* 2021. <https://doi.org/10.3390/cancers13194987>.
 107. Baeten KM, Akassoglou K. Extracellular matrix and matrix receptors in blood-brain barrier formation and stroke. *Dev Neurobiol.* 2011;71(11):1018–39.
 108. Aman J, Margadant C. Integrin-dependent cell-matrix adhesion in endothelial health and disease. *Circ Res.* 2023;132(3):355–78.
 109. Nguyen B, Bix G, Yao Y. Basal lamina changes in neurodegenerative disorders. *Mol Neurodegener.* 2021;16(1):81.
 110. Osada T, Gu YH, Kanazawa M, Tsubota Y, Hawkins BT, Spatz M, et al. Interendothelial claudin-5 expression depends on cerebral endothelial cell-matrix adhesion by β (1)-integrins. *J Cereb Blood Flow Metab.* 2011;31(10):1972–85.
 111. Kohn JC, Zhou DW, Bordeleau F, Zhou A, Mason B, Mitchell M, et al. Cooperative effects of matrix stiffness and fluid shear stress on endothelial cell behavior. *Biophys J.* 2015;108(3):471–8.
 112. Xue C, Zhang T, Xie X, Zhang Q, Zhang S, Zhu B, et al. Substrate stiffness regulates arterial-venous differentiation of endothelial progenitor cells via the Ras/Mek pathway. *Biochim et Biophys Acta (BBA) - Mol Cell Res.* 2017;1864(10):1799–808.
 113. Birukov KG, Birukova AA, Dudek SM, Verin AD, Crow MT, Zhan X, et al. Shear stress-mediated cytoskeletal remodeling and cortactin translocation in pulmonary endothelial cells. *Am J Respir Cell Mol Biol.* 2002;26(4):453–64.
 114. Uspenskaia O, Liebetrau M, Herms J, Danek A, Hamann GF. Aging is associated with increased collagen type IV accumulation in the basal lamina of human cerebral microvessels. *BMC Neurosci.* 2004;5:37.
 115. Hawkes CA, Gatherer M, Sharp MM, Dorr A, Yuen HM, Kalaria R, et al. Regional differences in the morphological and functional effects of aging on cerebral basement membranes and perivascular drainage of amyloid- β from the mouse brain. *Aging Cell.* 2013;12(2):224–36.
 116. Reed MJ, Damodarasamy M, Banks WA. The extracellular matrix of the blood-brain barrier: structural and functional roles in health, aging, and Alzheimer's disease. *Tissue Barriers.* 2019;7(4):1651157.
 117. Farkas E, De Jong GI, de Vos RAI, Jansen Steur ENH, Luiten PGM. Pathological features of cerebral cortical capillaries are doubled in Alzheimer's disease and Parkinson's disease. *Acta Neuropathol.* 2000;100(4):395–402.
 118. Mancardi GL, Perdelli F, Rivano C, Leonardi A, Bugiani O. Thickening of the basement membrane of cortical capillaries in Alzheimer's disease. *Acta Neuropathol.* 1980;49(1):79–83.
 119. Heinz A. Elastic fibers during aging and disease. *Ageing Res Rev.* 2021;66:101255.
 120. Fonck E, Feigl GG, Fasel J, Sage D, Unser M, Rüfenacht DA, Stergiopoulos N. Effect of aging on elastin functionality in human cerebral arteries. *Stroke.* 2009;40(7):2552–6.
 121. Girouard H, Iadecola C. Neurovascular coupling in the normal brain and in hypertension, stroke, and Alzheimer disease. *J Appl Physiol.* 2006;100(1):328–35.

122. Mandalà M, Cipolla MJ. Aging-related structural and functional changes in cerebral arteries: caloric restriction (CR) intervention. *J Vasc Med Surg*. 2021. <https://doi.org/10.1126/science.aav9518>.
123. Nortley R, Korte N, Izquierdo P, Hirunpattarasilp C, Mishra A, Jaunmuktane Z, et al. Amyloid β oligomers constrict human capillaries in Alzheimer's disease via signaling to pericytes. *Science*. 2019. <https://doi.org/10.1126/science.aav9518>.
124. Gürlér G, Soylu KO, Yemisci M. Importance of pericytes in the pathophysiology of cerebral ischemia. *Noro Psikiyatr Ars*. 2022;59(Suppl 1):S29–S35.
125. Verzijl N, DeGroot J, Thorpe SR, Bank RA, Shaw JN, Lyons TJ, et al. Effect of collagen turnover on the accumulation of advanced glycation end products*. *J Biol Chem*. 2000;275(50):39027–31.
126. Singh J, Di Ferrante N, Gyorkey F, Wilson N. Plasma glycosaminoglycans in normal individuals of various age. *Atherosclerosis*. 1977;28(3):319–24.
127. Libby JR, Royce H, Walker SR, Li L. The role of extracellular matrix in angiogenesis: beyond adhesion and structure. *Biomater Biosyst*. 2024;15:100097.
128. Lepelletier FX, Mann D, Robinson A, Pinteaux E, Boutin H. Early changes in extracellular matrix in Alzheimer's disease. *Neuropathol Appl Neurobiol*. 2017;43(2):167–82.
129. Tian J, Shi J, Smallman R, Iwatsubo T, Mann DMA. Relationships in Alzheimer's disease between the extent of A β deposition in cerebral blood vessel walls, as cerebral amyloid angiopathy, and the amount of cerebrovascular smooth muscle cells and collagen. *Neuropathol Appl Neurobiol*. 2006;32(3):332–40.
130. Benedict RH, DeLuca J, Phillips G, LaRocca N, Hudson LD, Rudick R. Validity of the symbol digit modalities test as a cognition performance outcome measure for multiple sclerosis. *Mult Scler*. 2017;23(5):721–33.
131. Zheng P, Pilutti LA, DuBose NG, Motl RW. Vascular function and cognition in persons with multiple sclerosis: preliminary examination. *Multiple Scler Related Disorders*. 2023;71:104578.
132. Bae JH, Kim JM, Park KY, Han SH. Association between arterial stiffness and the presence of cerebral small vessel disease markers. *Brain Behav*. 2021;11(1):e01935.
133. Miyagi T, Ishida A, Shinzato T, Ohya Y. Arterial stiffness is associated with small vessel disease irrespective of blood pressure in stroke-free individuals. *Stroke*. 2023;54(11):2814–21.
134. Boshra H, Awad M, Hussein M, Elyamani E. Vascular dysfunction and dyslipidemia in multiple sclerosis: are they correlated with disease duration and disability status? *Egypt Heart J*. 2022;74(1):9.
135. Alvarez-Bueno C, Cunha PG, Martinez-Vizcaino V, Pozuelo-Carrascosa DP, Visier-Alfonso ME, Jimenez-Lopez E, Cervero-Redondo I. Arterial stiffness and cognition among adults: a systematic review and meta-analysis of observational and longitudinal studies. *J Am Heart Assoc*. 2020;9(5):e014621.
136. Ohashi KL, Tung DK, Wilson J, Zweifach BW, Schmid-Schönbein GW. Transvascular and interstitial migration of neutrophils in rat mesentery. *Microcirculation*. 1996;3(2):199–210.
137. Yadav R, Larbi KY, Young RE, Nourshargh S. Migration of leukocytes through the vessel wall and beyond. *Thromb Haemost*. 2003;90(4):598–606.
138. Bartholomäus I, Kawakami N, Odoardi F, Schläger C, Miljkovic D, Ellwart JW, et al. Effector T cell interactions with meningeal vascular structures in nascent autoimmune CNS lesions. *Nature*. 2009;462(7269):94–8.
139. Zhang X, Wang Y, Song J, Gerwien H, Chuquisana O, Chashchina A, et al. The endothelial basement membrane acts as a checkpoint for entry of pathogenic T cells into the brain. *J Exp Med*. 2020. <https://doi.org/10.1084/jem.20191339>.
140. Riddle RB, Jennbacken K, Hansson KM, Harper MT. Endothelial inflammation and neutrophil transmigration are modulated by extracellular matrix composition in an inflammation-on-a-chip model. *Sci Rep*. 2022;12(1):6855.
141. Mutgan AC, Jandl K, Kwapiszewska G. Endothelial basement membrane components and their products, matrikines: active drivers of pulmonary hypertension? *Cells*. 2020. <https://doi.org/10.3390/cells9092029>.
142. Leclech C, Natale CF, Barakat AI. The basement membrane as a structured surface—role in vascular health and disease. *J Cell Sci*. 2020. <https://doi.org/10.1242/jcs.239889>.
143. Engelhardt B. Regulation of immune cell entry into the central nervous system. *Results Probl Cell Differ*. 2006;43:259–80.
144. Haage V, De Jager PL. Neuroimmune contributions to Alzheimer's disease: a focus on human data. *Mol Psychiatry*. 2022;27(8):3164–81.
145. Merlini M, Kirabali T, Kulic L, Nitsch RM, Ferretti MT. Extravascular CD3+ T cells in brains of Alzheimer disease patients correlate with tau but not with amyloid pathology: an immunohistochemical study. *Neurodegener Dis*. 2018;18(1):49–56.
146. Zeng J, Liao Z, Yang H, Wang Q, Wu Z, Hua F, Zhou Z. T cell infiltration mediates neurodegeneration and cognitive decline in Alzheimer's disease. *Neurobiol Dis*. 2024;193:106461.
147. Rossy J, Laufer JM, Legler DF. Role of mechanotransduction and tension in T cell function. *Front Immunol*. 2018. <https://doi.org/10.3389/fimmu.2018.02638>.
148. Schaefer A, Hordijk PL. Cell-stiffness-induced mechanosignaling—a key driver of leukocyte transendothelial migration. *J Cell Sci*. 2015;128(13):2221–30.
149. Sun C, Migliorini C, Munn LL. Red blood cells initiate leukocyte rolling in postcapillary expansions: a lattice boltzmann analysis. *Biophys J*. 2003;85(1):208–22.
150. Muller WA. Getting leukocytes to the site of inflammation. *Vet Pathol*. 2013;50(1):7–22.
151. Zhang X, Chen A, De Leon D, Li H, Noiri E, Moy VT, Goligorsky MS. Atomic force microscopy measurement of leukocyte-endothelial interaction. *Am J Physiol Heart Circ Physiol*. 2004;286(1):H359–67.
152. Engelhardt B, Ransohoff RM. Capture, crawl, cross: the T cell code to breach the blood-brain barriers. *Trends Immunol*. 2012;33(12):579–89.
153. Siegelman MH, Stanescu D, Estess P. The CD44-initiated pathway of T-cell extravasation uses VLA-4 but not LFA-1 for firm adhesion. *J Clin Invest*. 2000;105(5):683–91.
154. Coisne C, Favéu C, Delplace Y, Dehouck L, Miller F, Cecchelli R, Dehouck B. Differential expression of selectins by mouse brain capillary endothelial cells in vitro in response to distinct inflammatory stimuli. *Neurosci Lett*. 2006;392(3):216–20.
155. McDonnell GV, McMillan SA, Douglas JP, Droogan AG, Hawkins SA. Serum soluble adhesion molecules in multiple sclerosis: raised sVCAM-1, sICAM-1 and sE-selectin in primary progressive disease. *J Neurol*. 1999;246(2):87–92.
156. Yousef H, Czupalla CJ, Lee D, Chen MB, Burke AN, Zera KA, et al. Aged blood impairs hippocampal neural precursor activity and activates microglia via brain endothelial cell VCAM1. *Nat Med*. 2019;25(6):988–1000.
157. Sathiyaradan K, Coisne C, Enzmann G, Deutsch U, Engelhardt B. PSGL-1 and E/P-selectins are essential for T-cell rolling in inflamed CNS microvessels but dispensable for initiation of EAE. *Eur J Immunol*. 2014;44(8):2287–94.
158. Nishihara H, Soldati S, Mossu A, Rosito M, Rudolph H, Muller WA, et al. Human CD4+ T cell subsets differ in their abilities to cross endothelial and epithelial brain barriers in vitro. *Fluids and Barriers of the CNS*. 2020;17(1):3.
159. Mapunda JA, Tibar H, Regragui W, Engelhardt B. How does the immune system enter the brain? *Front Immunol*. 2022. <https://doi.org/10.3389/fimmu.2022.805657/full>.
160. Lyck R, Engelhardt B. Going against the tide—how encephalitogenic T cells breach the blood-brain barrier. *J Vasc Res*. 2012;49(6):497–509.
161. Steiner O, Coisne C, Cecchelli R, Boscacci R, Deutsch U, Engelhardt B, Lyck R. Differential roles for endothelial ICAM-1, ICAM-2, and VCAM-1 in shear-resistant T cell arrest, polarization, and directed crawling on blood-brain barrier endothelium. *J Immunol*. 2010;185(8):4846–55.
162. Zhu C, Chen W, Lou J, Rittase W, Li K. Mechanosensing through immunoreceptors. *Nat Immunol*. 2019;20(10):1269–78.
163. Pigeon SV, Govendir MA, Kempe D, Biro M. Mechanoimmunology: molecular-scale forces govern immune cell functions. *Mol Biol Cell*. 2018;29(16):1919–26.
164. Liu B, Chen W, Zhu C. Accumulation of dynamic catch bonds between TCR and agonist peptide-MHC triggers T cell signaling. *Cell*. 2014;157(2):357–68.
165. Harrison DL, Fang Y, Huang J. T-cell mechanobiology: force sensation, potentiation, and translation. *Front Phys*. 2019. <https://doi.org/10.3389/fphys.2019.00045/full>.
166. Alon R, Dustin ML. Force as a facilitator of integrin conformational changes during leukocyte arrest on blood vessels and antigen-presenting cells. *Immunity*. 2007;26(1):17–27.

167. Sun Z, Costell M, Fässler R. Integrin activation by talin, kindlin and mechanical forces. *Nat Cell Biol.* 2019;21(1):25–31.
168. Zhu J, Luo B-H, Xiao T, Zhang C, Nishida N, Springer TA. Structure of a complete integrin ectodomain in a physiologic resting state and activation and deactivation by applied forces. *Mol Cell.* 2008;32(6):849–61.
169. Shen B, Delaney MK, Du X. Inside-out, outside-in, and inside-outside-in: G protein signaling in integrin-mediated cell adhesion, spreading, and retraction. *Curr Opin Cell Biol.* 2012;24(5):600–6.
170. Shao B, Yago T, Panicker SR, Zhang N, Liu Z, McEver RP. Th1 cells rolling on selectins trigger DAP12-dependent signals that activate integrin $\alpha\text{L}\beta\text{2}$. *J Immunol.* 2020;204(1):37–48.
171. Woolf E, Grigorova I, Sagiv A, Grabovsky V, Feigelson SW, Shulman Z, et al. Lymph node chemokines promote sustained T lymphocyte motility without triggering stable integrin adhesiveness in the absence of shear forces. *Nat Immunol.* 2007;8(10):1076–85.
172. Dominguez GA, Anderson NR, Hammer DA. The direction of migration of T-lymphocytes under flow depends upon which adhesion receptors are engaged. *Integr Biol.* 2015;7(3):345–55.
173. Lee J, Song KH, Kim T, Doh J. Endothelial cell focal adhesion regulates transendothelial migration and subendothelial crawling of T cells. *Front Immunol.* 2018. <https://doi.org/10.3389/fimmu.2018.00048/full>.
174. Coisne C, Lyck R, Engelhardt B. Live cell imaging techniques to study T cell trafficking across the blood-brain barrier in vitro and in vivo. *Fluids Barriers CNS.* 2013;10(1):7.
175. Elosegui-Artola A, Oria R, Chen Y, Kosmalska A, Pérez-González C, Castro N, et al. Mechanical regulation of a molecular clutch defines force transmission and transduction in response to matrix rigidity. *Nat Cell Biol.* 2016;18(5):540–8.
176. Renkawitz J, Sixt M. Mechanisms of force generation and force transmission during interstitial leukocyte migration. *EMBO Rep.* 2010;11(10):744–50.
177. Saitakis M, Dogniaux S, Goudot C, Bui N, Asnacios S, Maurin M, et al. Different TCR-induced T lymphocyte responses are potentiated by stiffness with variable sensitivity. *Elife.* 2017. <https://doi.org/10.7554/elife.23190>.
178. Schaefer A, Te Riet J, Ritz K, Hoogenboezem M, Anthony EC, Mul FPJ, et al. Actin-binding proteins differentially regulate endothelial cell stiffness, ICAM-1 function and neutrophil transmigration. *J Cell Sci.* 2014;127(20):4470–82.
179. Sumagin R, Prizant H, Lomakina E, Waugh RE, Sarelis IH. LFA-1 and Mac-1 define characteristically different intraluminal crawling and emigration patterns for monocytes and neutrophils in situ. *J Immunol.* 2010;185(11):7057–66.
180. Kim SHJ, Hammer DA. Integrin crosstalk allows CD4⁺ T lymphocytes to continue migrating in the upstream direction after flow. *Integr Biol.* 2019;11(10):384–93.
181. Schimmel L, Heemskerk N, van Buul JD. Leukocyte transendothelial migration: a local affair. *Small GTPases.* 2017;8(1):1–15.
182. Martinelli R, Zeiger A, Whitfield M, Sciuto T, Dvorak A, Van Vliet K, et al. Probing the biomechanical contribution of the endothelium to lymphocyte migration: diapedesis by the path of least resistance. *J Cell Sci.* 2014. <https://doi.org/10.1242/jcs.148619>.
183. Abadier M, Haghighyegh Jahromi N, Cardoso Alves L, Boscacci R, Vestweber D, Barnum S, et al. Cell surface levels of endothelial ICAM-1 influence the transcellular or paracellular T-cell diapedesis across the blood-brain barrier. *Eur J Immunol.* 2015;45(4):1043–58.
184. Marchetti L, Francisco D, Soldati S, Haghighyegh Jahromi N, Barcos S, Gruber I, et al. ACKR1 favors transcellular over paracellular T-cell diapedesis across the blood-brain barrier in neuroinflammation in vitro. *Eur J Immunol.* 2022;52(1):161–77.
185. Lutz SE, Smith JR, Kim DH, Olson CVL, Ellefsen K, Bates JM, et al. Caveolin1 is required for Th1 cell infiltration, but not tight junction remodeling, at the blood-brain barrier in autoimmune neuroinflammation. *Cell Rep.* 2017;21(8):2104–17.
186. Trevino TN, Almousawi AA, Ochoa-Raya A, Zemanski K, Oliveira SD, Marottoli FM, et al. Endothelial caveolin-1 and CXCL10 promote transcellular migration of autoreactive T cells across the blood-brain barrier. *bioRxiv.* 2022. <https://doi.org/10.1101/2022.11.15.516689v1.abstract>.
187. Bennett J, Basivreddy J, Kollar A, Biron KE, Reickmann P, Jefferies WA, McQuaid S. Blood–brain barrier disruption and enhanced vascular permeability in the multiple sclerosis model EAE. *J Neuroimmunol.* 2010;229(1):180–91.
188. Carman CV. Mechanisms for transcellular diapedesis: probing and pathfinding by ‘invadosome-like protrusions’. *J Cell Sci.* 2009;122(17):3025–35.
189. Alon R, Van Buul JD. Leukocyte breaching of endothelial barriers: the actin link. *Trends Immunol.* 2017;38(8):606–15.
190. Carman CV, Sage PT, Sciuto TE, De La Fuente MA, Geha RS, et al. Transcellular diapedesis is initiated by invasive podosomes. *Immunity.* 2007;26(6):784–97.
191. Heemskerk N, Schimmel L, Oort C, Van Rijssel J, Yin T, Ma B, et al. F-actin-rich contractile endothelial pores prevent vascular leakage during leukocyte diapedesis through local RhoA signalling. *Nat Commun.* 2016;7(1):10493.
192. Smyth LA, Afzali B, Tsang J, Lombardi G, Lechler RI. Intercellular transfer of MHC and immunological molecules: molecular mechanisms and biological significance. *Am J Transplant.* 2007;7(6):1442–9.
193. Brezinschek RI, Oppenheimer-Marks N, Lipsky PE. Activated T cells acquire endothelial cell surface determinants during transendothelial migration. *J Immunol.* 1999;162(3):1677–84.
194. Grönloh MLB, Arts JGG, Palacios Martínez S, van der Veen AA, Kempers L, van Steen ACI, et al. Endothelial transmigration hotspots limit vascular leakage through heterogeneous expression of ICAM-1. *EMBO Rep.* 2023;24(1):e55483.
195. Dorland YL, Huveneers S. Cell–cell junctional mechanotransduction in endothelial remodeling. *Cell Mol Life Sci.* 2017;74(2):279–92.
196. Twiss F, Le Duc Q, Van Der Horst S, Tabdili H, Van Der Krogt G, Wang N, et al. Vinculin-dependent cadherin mechanosensing regulates efficient epithelial barrier formation. *Biol Open.* 2012;1(11):1128–40.
197. Malinova TS, Angulo-Urarte A, Nüchel J, Tauber M, Van Der Stoep MM, Janssen V, et al. A junctional PACSIN2/EHD4/MICAL-L1 complex coordinates VE-cadherin trafficking for endothelial migration and angiogenesis. *Nature Communications.* 2021;12(1).
198. Arbore C, Sergides M, Gardini L, Bianchi G, Kashchuk AV, Pertici I, et al. α -catenin switches between a slip and an asymmetric catch bond with F-actin to cooperatively regulate cell junction fluidity. *bioRxiv.* 2021. <https://doi.org/10.1038/s41467-022-28779-7>.
199. Huang DL, Bax NA, Buckley CD, Weis WI, Dunn AR. Vinculin forms a directionally asymmetric catch bond with F-actin. *Science.* 2017;357(6352):703–6.
200. Owen LM, Bax NA, Weis WI, Dunn AR. The C-terminal actin-binding domain of talin forms an asymmetric catch bond with F-actin. *Proc Natl Acad Sci USA.* 2022;119(10):e2109329119.
201. Arbore C, Sergides M, Gardini L, Bianchi G, Kashchuk AV, Pertici I, et al. α -catenin switches between a slip and an asymmetric catch bond with F-actin to cooperatively regulate cell junction fluidity. *Nat Commun.* 2022;13(1):1146.
202. Fang Y, Wu D, Birukov KG. Mechanosensing and mechanoregulation of endothelial cell functions. *Compr Physiol.* 2019;9(2):873–904.
203. Dembo M, Torney DC, Saxman K, Hammer D, Murray JD. The reaction-limited kinetics of membrane-to-surface adhesion and detachment. *Proc R Soc Lond B.* 1988;234(1274):55–83.
204. Collins C, Guilluy C, Welch C, O'Brien ET, Hahn K, Superfine R, et al. Localized tensional forces on PECAM-1 elicit a global mechanotransduction response via the integrin-RhoA pathway. *Curr Biol.* 2012;22(22):2087–94.
205. Gavard J, Gutkind JS. VE-cadherin and claudin-5: it takes two to tango. *Nat Cell Biol.* 2008;10(8):883–5.
206. Wimmer I, Tietz S, Nishihara H, Deutsch U, Sallusto F, Gosselet F, et al. PECAM-1 stabilizes blood-brain barrier integrity and favors paracellular T-cell diapedesis across the blood-brain barrier during neuroinflammation. *Front Immunol.* 2019. <https://doi.org/10.3389/fimmu.2019.00711>.
207. Tzima E, Irani-Tehrani M, Kiosses WB, Dejana E, Schultz DA, Engelhardt B, et al. A mechanosensory complex that mediates the endothelial cell response to fluid shear stress. *Nature.* 2005;437(7057):426–31.
208. Collins C, Guilluy C, Welch CE, Hahn K, Superfine R, et al. Localized tensional forces on PECAM-1 elicit a global mechanotransduction response via the integrin-RhoA pathway. *Curr Biol.* 2012;22(22):2087–94.
209. Weber EW, Han F, Tauseef M, Birnbaumer L, Mehta D, Muller WA. TRPC6 is the endothelial calcium channel that regulates leukocyte transendothelial migration during the inflammatory response. *J Exp Med.* 2015;212(11):1883–99.

210. Winneberger J, Schöls S, Lessmann K, Rándež-Garbayo J, Bauer AT, Mohamud Yusuf A, et al. Platelet endothelial cell adhesion molecule-1 is a gatekeeper of neutrophil transendothelial migration in ischemic stroke. *Brain Behav Immun*. 2021;93:277–87.
211. Harry BL, Sanders JM, Feaver RE, Lansey M, Deem TL, Zarbock A, et al. Endothelial cell PECAM-1 promotes atherosclerotic lesions in areas of disturbed flow in ApoE-deficient mice. *Arterioscler Thromb Vasc Biol*. 2008;28(11):2003–8.
212. Haas AJ, Zihni C, Ruppel A, Hartmann C, Ebnet K, Tada M, et al. Interplay between extracellular matrix stiffness and JAM-A regulates mechanical load on ZO-1 and tight junction assembly. *Cell Rep*. 2020;32(3):107924.
213. Chistiakov DA, Orekhov AN, Bobryshev YV. Effects of shear stress on endothelial cells: go with the flow. *Acta Physiol*. 2017;219(2):382–408.
214. Tornavaca O, Chia M, Dufton N, Almagro LO, Conway DE, Randi AM, et al. ZO-1 controls endothelial adherens junctions, cell–cell tension, angiogenesis, and barrier formation. *J Cell Biol*. 2015;208(6):821–38.
215. Liu Z, Tan JL, Cohen DM, Yang MT, Sniadecki NJ, Ruiz SA, et al. Mechanical tugging force regulates the size of cell-cell junctions. *Proc Natl Acad Sci*. 2010;107(22):9944–9.
216. Yap AS, Duszyc K, Viasnoff V. Mechanosensing and mechanotransduction at cell-cell junctions. *Cold Spring Harb Perspect Biol*. 2018. <https://doi.org/10.1101/cshperspect.a028761>.
217. Shihata W, Michell D, Andrews K, Chin-Dusting J. Caveolae: a role in endothelial inflammation and mechanotransduction? *Front Physiol*. 2016. <https://doi.org/10.3389/fphys.2016.00628>.
218. Jin P, Jan LY, Jan Y-N. Mechanosensitive ion channels: structural features relevant to mechanotransduction mechanisms. *Ann Rev Neurosci*. 2020;43:207–29.
219. Arnadóttir J, Chalfie M. Eukaryotic mechanosensitive channels. *Annu Rev Biophys*. 2010;39:111–37.
220. Beedle AEM, Roca-Cusachs P. The reversibility of cellular mechanotransduction. *Curr Opin Cell Biol*. 2023;84:102229.
221. Sala S, Caillier A, Oakes PW. Principles and regulation of mechanosensing. *J Cell Sci*. 2024. <https://doi.org/10.1242/jcs.261338>.
222. Jin Z-G, Ueba H, Tanimoto T, Lungu AO, Frame MD, Berk BC. Ligand-independent activation of vascular endothelial growth factor receptor 2 by fluid shear stress regulates activation of endothelial nitric oxide synthase. *Circ Res*. 2003;93(4):354–63.
223. Jong Lee H, Young KG. Shear stress activates Tie2 receptor tyrosine kinase in human endothelial cells. *Biochem Biophys Res Commun*. 2003;304(2):399–404.
224. Gavard J, Gutkind JS. VEGF controls endothelial-cell permeability by promoting the beta-arrestin-dependent endocytosis of VE-cadherin. *Nat Cell Biol*. 2006;8(11):1223–34.
225. Miller B, Sewell-Loftin MK. Mechanoregulation of vascular endothelial growth factor receptor 2 in angiogenesis. *Front Cardiovasc Med*. 2021;8:804934.
226. Matsuo E, Okamoto T, Ito A, Kawamoto E, Asanuma K, Wada K, et al. Substrate stiffness modulates endothelial cell function via the YAP-Dll4-Notch1 pathway. *Exp Cell Res*. 2021;408(1):112835.
227. Wang KC, Yeh YT, Nguyen P, Limquenco E, Lopez J, Thorossian S, et al. Flow-dependent YAP/TAZ activities regulate endothelial phenotypes and atherosclerosis. *Proc Natl Acad Sci USA*. 2016;113(41):11525–30.
228. Huang HC, Wang TY, Rousseau J, Mungeray M, Michaud C, Plaisier C, et al. Lesion-specific suppression of YAP/TAZ by biomimetic nanodrug ameliorates atherosclerosis development. *bioRxiv*. 2023;16:389.
229. Jia G, Aroor AR, Jia C, Sowers JR. Endothelial cell senescence in aging-related vascular dysfunction. *Biochim Biophys Acta Mol Basis Dis*. 2019;1865(7):1802–9.
230. Chachivili M, Zhang YL, Frangos JA. G protein-coupled receptors sense fluid shear stress in endothelial cells. *Proc Natl Acad Sci*. 2006;103(42):15463–8.
231. Xu J, Mathur J, Vessières E, Hammack S, Nonomura K, Favre J, et al. GPR68 senses flow and is essential for vascular physiology. *Cell*. 2018;173(3):762–75.e16.
232. Iliff AJ, Xu XZ. A mechanosensitive GPCR that detects the bloody force. *Cell*. 2018;173(3):542–4.
233. Ludwig M-G, Vanek M, Guerini D, Gasser JA, Jones CE, Junker U, et al. Proton-sensing G-protein-coupled receptors. *Nature*. 2003;425(6953):93–8.
234. Wenzel J, Hansen CE, Bettoni C, Vogt MA, Lembrich B, Natsagdorj R, et al. Impaired endothelium-mediated cerebrovascular reactivity promotes anxiety and respiration disorders in mice. *Proc Natl Acad Sci*. 2020;117(3):1753–61.
235. Wang T, Zhou G, He M, Xu Y, Rusyniak WG, Xu Y, et al. GPR68 is a neuroprotective proton receptor in brain ischemia. *Stroke*. 2020;51(12):3690–700.
236. McGinley MP, Cohen JA. Sphingosine 1-phosphate receptor modulators in multiple sclerosis and other conditions. *Lancet*. 2021;398(10306):1184–94.
237. Jung B, Obinata H, Galvani S, Mendelson K, Ding B-S, Skoura A, et al. Flow-regulated endothelial S1P receptor-1 signaling sustains vascular development. *Dev Cell*. 2012;23(3):600–10.
238. Fu Y, Hao J, Zhang N, Ren L, Sun N, Li YJ, et al. Fingolimod for the treatment of intracerebral hemorrhage: a 2-arm proof-of-concept study. *JAMA Neurol*. 2014;71(9):1092–101.
239. Zhu Z, Fu Y, Tian D, Sun N, Han W, Chang G, et al. Combination of the immune modulator fingolimod with alteplase in acute ischemic stroke: a pilot trial. *Circulation*. 2015;132(12):1104–12.
240. Nishihara H, Shimizu F, Sano Y, Takeshita Y, Maeda T, Abe M, et al. Fingolimod prevents blood-brain barrier disruption induced by the sera from patients with multiple sclerosis. *PLoS ONE*. 2015;10(3):e0121488.
241. Cantalupo A, Gargiulo A, Dautaj E, Liu C, Zhang Y, Hla T, Di Lorenzo A. S1PR1 (sphingosine-1-phosphate receptor 1) signaling regulates blood flow and pressure. *Hypertension*. 2017;70(2):426–34.
242. Hughes R, Dalakas MC, Merkies I, Latov N, Léger J-M, Nobile-Orazio E, et al. Oral fingolimod for chronic inflammatory demyelinating polyradiculoneuropathy (FORCIP Trial): a double-blind, multicentre, randomised controlled trial. *Lancet Neurol*. 2018;17(8):689–98.
243. Schwab A, Fabian A, Hanley PJ, Stock C. Role of ion channels and transporters in cell migration. *Physiol Rev*. 2012;92(4):1865–913.
244. Tanaka K, Joshi D, Timalina S, Schwartz MA. Early events in endothelial flow sensing. *Cytoskeleton*. 2021;78(6):217–31.
245. Ma X, Liu W. Calcium signaling in brain microvascular endothelial cells and its roles in the function of the blood-brain barrier. *NeuroReport*. 2019;30(18):1271–7.
246. Brown RC, Wu L, Hicks K, O'Neil RG. Regulation of blood-brain barrier permeability by transient receptor potential type C and type v calcium-permeable channels. *Microcirculation*. 2008;15(4):359–71.
247. Berrout J, Jin M, O'Neil RG. Critical role of TRPP2 and TRPC1 channels in stretch-induced injury of blood-brain barrier endothelial cells. *Brain Res*. 2012;1436:1–12.
248. Hansen CE, Kamermans A, Mol K, Berve K, Rodriguez-Mogeda C, Fung WK, et al. Inflammation-induced TRPV4 channels exacerbate blood-brain barrier dysfunction in multiple sclerosis. *J Neuroinflammation*. 2024;21(1):72.
249. Li J, Hou B, Tumova S, Muraki K, Bruns A, Ludlow MJ, et al. Piezo1 integration of vascular architecture with physiological force. *Nature*. 2014;515(7526):279–82.
250. Harraz OF, Klug NR, Senatore AJ, Hill-Eubanks DC, Nelson MT. Piezo1 is a mechanosensor channel in central nervous system capillaries. *Circ Res*. 2022;130(10):1531–46.
251. Kang H, Hong Z, Zhong M, Klomp J, Bayless KJ, Mehta D, et al. Piezo1 mediates angiogenesis through activation of MT1-MMP signaling. *Am J Phys Cell Physiol*. 2019;316(1):C92–103.
252. Friedrich EE, Hong Z, Xiong S, Zhong M, Di A, Rehman J, et al. Endothelial cell Piezo1 mediates pressure-induced lung vascular hyperpermeability via disruption of adherens junctions. *Proc Natl Acad Sci USA*. 2019;116(26):12980–5.
253. Wang Z, Chen J, Babicheva A, Jain PP, Rodriguez M, Ayon RJ, et al. Endothelial upregulation of mechanosensitive channel Piezo1 in pulmonary hypertension. *Am J Physiol Cell Physiol*. 2021;321(6):C1010–27.
254. Yang Y, Wang D, Zhang C, Yang W, Li C, Gao Z, et al. Piezo1 mediates endothelial atherogenic inflammatory responses via regulation of YAP/TAZ activation. *Hum Cell*. 2021. <https://doi.org/10.1007/s13577-021-00600-5>.
255. Albarrán-Juárez J, Iring A, Wang S, Joseph S, Grimm M, Strilic B, et al. Piezo1 and Gq/G11 promote endothelial inflammation depending on flow pattern and integrin activation. *J Exp Med*. 2018;215(10):2655–72.

256. Chuntharpursat-Bon E, Povstyan OV, Ludlow MJ, Carrier DJ, Debant M, Shi J, et al. PIEZO1 and PECAM1 interact at cell-cell junctions and partner in endothelial force sensing. *Commun Biol*. 2023;6(1):358.
257. Bertin S, Raz E. Transient receptor potential (TRP) channels in T cells. *Semin Immunopathol*. 2016;38(3):309–19.
258. Rosenbaum T, Islas LD. Molecular physiology of TRPV channels: controversies and future challenges. *Annu Rev Physiol*. 2023;85:293–316.
259. Nikolaev YA, Cox CD, Ridone P, Rohde PR, Cordero-Morales JF, Vásquez V, et al. Mammalian TRP ion channels are insensitive to membrane stretch. *J Cell Sci*. 2019;132(23):238360.
260. Hill-Eubanks DC, Gonzales AL, Sonkusare SK, Nelson MT. Vascular TRP channels: performing under pressure and going with the flow. *Physiology*. 2014;29(5):343–60.
261. Dragovich MA, Chester D, Fu BM, Wu C, Xu Y, Goligorsky MS, Zhang XF. Mechanotransduction of the endothelial glycocalyx mediates nitric oxide production through activation of TRP channels. *Am J Physiol Cell Physiol*. 2016;311(6):C846–53.
262. Earley S, Gonzales AL, Crnich R. Endothelium-dependent cerebral artery dilation mediated by TRPA1 and Ca²⁺-activated K⁺ channels. *Circ Res*. 2009;104(8):987–94.
263. Tiwari A, Acharya T, Saha S, et al. Transient receptor potential ankyrin1 channel is endogenously expressed in T cells and is involved in immune functions. *Biosci Rep*. 2019;39(9):20191437.
264. Naert R, López-Requena A, Talavera K. TRPA1 expression and pathophysiology in immune cells. *Int J Mol Sci*. 2021;22(21):11460.
265. Wenning AS, Neblung K, Strauß B, Wolfs M-J, Sappok A, Hoth M, Schwarz EC. TRP expression pattern and the functional importance of TRPC3 in primary human T-cells. *Biochim et Biophys Acta (BBA) - Mol Cell Res*. 2011;1813(3):412–23.
266. Yip H, Chan W-Y, Leung P-C, Kwan H-Y, Liu C, Huang Y, et al. Expression of TRPC homologs in endothelial cells and smooth muscle layers of human arteries. *Histochem Cell Biol*. 2004;122(6):553–61.
267. Lindemann O, Strodthoff C, Horstmann M, Nielsen N, Jung F, Schimmelpfennig S, et al. TRPC1 regulates fMLP-stimulated migration and chemotaxis of neutrophil granulocytes. *Biochim et Biophys Acta (BBA) - Mol Cell Res*. 2015;1853(9):2122–30.
268. Chauhan A, Sun Y, Sukumaran P, Quenum Zangbede FO, Jondle CN, Sharma A, et al. M1 macrophage polarization is dependent on TRPC1-mediated calcium entry. *iScience*. 2018;8:85–102.
269. Brécard S, Melchior C, Plançon S, Schenten V, Tschirhart EJ. Store-operated Ca²⁺ channels formed by TRPC1, TRPC6 and orai1 and non-store-operated channels formed by TRPC3 are involved in the regulation of NADPH oxidase in HL-60 granulocytes. *Cell Calcium*. 2008;44(5):492–506.
270. Tano JY, Solanki S, Lee RH, Smedlund K, Birnbaumer L, Vazquez G. Bone marrow deficiency of TRPC3 channel reduces early lesion burden and necrotic core of advanced plaques in a mouse model of atherosclerosis. *Cardiovasc Res*. 2014;101(1):138–44.
271. Numaga T, Nishida M, Kiyonaka S, Kato K, Katano M, Mori E, et al. Ca²⁺ influx and protein scaffolding via TRPC3 sustain PKC β and ERK activation in B cells. *J Cell Sci*. 2010;123(Pt 6):927–38.
272. Quick K, Zhao J, Eijkelkamp N, Linley JE, Rugiero F, Cox JJ, et al. TRPC3 and TRPC6 are essential for normal mechanotransduction in subsets of sensory neurons and cochlear hair cells. *Open Biol*. 2012;2(5):120068.
273. Tao L, Guo G, Qi Y, Xiong Y, Ma X, Wu N, et al. Inhibition of canonical transient receptor potential 5 channels polarizes macrophages to an M1 phenotype. *Pharmacology*. 2020;105(3–4):202–8.
274. Shen B, Wong C-O, Lau O-C, Woo T, Bai S, Huang Y, Yao X. Plasma membrane mechanical stress activates TRPC5 channels. *PLoS ONE*. 2015;10(4):e0122227.
275. Lindemann O, Rossaint J, Najder K, Schimmelpfennig S, Hofschroer V, Wälte M, et al. Intravascular adhesion and recruitment of neutrophils in response to CXCL1 depends on their TRPC6 channels. *J Mol Med*. 2020;98(3):349–60.
276. Liu Z, Yang J, Zhang X, Xu P, Zhang T, Yang Z. Developmental changes in the expression and function of TRPC6 channels related the F-actin organization during differentiation in podocytes. *Cell Calcium*. 2015;58(6):541–8.
277. Riazanski V, Gabdoulkhakova AG, Boynton LS, Eguchi RR, Deriy LV, Hogarth DK, et al. TRPC6 channel translocation into phagosomal membrane augments phagosomal function. *Proc Natl Acad Sci*. 2015;112(47):E6486–95.
278. Spassova MA, Hewavitharana T, Xu W, Soboloff J, Gill DL. A common mechanism underlies stretch activation and receptor activation of TRPC6 channels. *Proc Natl Acad Sci*. 2006;103(44):16586–91.
279. Serafini N, Dahdah A, Barbet G, Demion M, Attout T, Gautier G, et al. The TRPM4 channel controls monocyte and macrophage, but not neutrophil, function for survival in sepsis. *J Immunol*. 2012;189(7):3689–99.
280. Morita H, Honda A, Inoue R, Ito Y, Abe K, Nelson MT, Brayden JE. Membrane stretch-induced activation of a TRPM4-like nonselective cation channel in cerebral artery myocytes. *J Pharmacol Sci*. 2007;103(4):417–26.
281. Gerzanich V, Kwon MS, Woo SK, Ivanov A, Simard JM. SUR1-TRPM4 channel activation and phasic secretion of MMP-9 induced by tPA in brain endothelial cells. *PLoS ONE*. 2018;13(4):e0195526.
282. Son M-J, Kim J-C, Kim SW, Chidipi B, Muniyandi J, Singh TD, et al. Shear stress activates monovalent cation channel transient receptor potential melastatin subfamily 4 in rat atrial myocytes via type 2 inositol 1,4,5-trisphosphate receptors and Ca²⁺-release. *J Physiol*. 2016;594(11):2985–3004.
283. Zeng Z, Inoue K, Sun H, Leng T, Feng X, Zhu L, Xiong Z-G. TRPM7 regulates vascular endothelial cell adhesion and tube formation. *Am J Physiol Cell Physiol*. 2015;308(4):C308–18.
284. Nauli SM, Kawanabe Y, Kaminski JJ, Pearce WJ, Ingber DE, Zhou J. Endothelial cilia are fluid shear sensors that regulate calcium signaling and nitric oxide production through polycystin-1. *Circulation*. 2008;117(9):1161–71.
285. Majhi RK, Sahoo SS, Yadav M, Pratheek BM, Chattopadhyay S, Goswami C. Functional expression of TRPV channels in T cells and their implications in immune regulation. *FEBS J*. 2015;282(14):2661–81.
286. Luo H, Saubamea B, Chasseigneaux S, Cochois V, Smirnova M, Glacial F, et al. Molecular and functional study of transient receptor potential vanilloid 1–4 at the rat and human blood-brain barrier reveals interspecies differences. *Front Cell Dev Biol*. 2020. <https://doi.org/10.3389/fcell.2020.578514>.
287. Bertin S, Aoki-Nonaka Y, De Jong PR, Nohara LL, Xu H, Stanwood SR, et al. The ion channel TRPV1 regulates the activation and proinflammatory properties of CD4⁺ T cells. *Nat Immunol*. 2014;15(11):1055–63.
288. Heiner I, Eisfeld J, Halaszovich CR, Wehage E, Jüngling E, Zitt C, Lückhoff A. Expression profile of the transient receptor potential (TRP) family in neutrophil granulocytes: evidence for currents through long TRP channel 2 induced by ADP-ribose and NAD. *Biochem J*. 2003;371(3):1045–53.
289. Pottosin I, Delgado-Enciso I, Bonales-Alatorre E, Nieto-Pescador MG, Moreno-Galindo EG, Dobrovinskaya O. Mechanosensitive Ca²⁺-permeable channels in human leukemic cells: pharmacological and molecular evidence for TRPV2. *Biochim Biophys Acta (BBA) - Biomembranes*. 2015;1848(1):51–9.
290. Nagasawa M, Kojima I. Translocation of TRPV2 channel induced by focal administration of mechanical stress. *Physiol Rep*. 2015. <https://doi.org/10.1481/phy2.12296>.
291. Marrelli SP, O'Neil RG, Brown RC, Bryan RM. PLA2 and TRPV4 channels regulate endothelial calcium in cerebral arteries. *Am J Physiol-Heart Circ Physiol*. 2007;292(3):H1390–7.
292. Matthews BD, Thodeti CK, Tytell JD, Mammoto A, Overby DR, Ingber DE. Ultra-rapid activation of TRPV4 ion channels by mechanical forces applied to cell surface β 1 integrins. *Integr Biol*. 2010;2(9):435.
293. Vriens J, Watanabe H, Janssens A, Droogmans G, Voets T, Nilius B. Cell swelling, heat, and chemical agonists use distinct pathways for the activation of the cation channel TRPV4. *Proc Natl Acad Sci*. 2004;101(1):396–401.
294. Yin J, Michalick L, Tang C, Tabuchi A, Goldenberg N, Dan Q, et al. Role of transient receptor potential vanilloid 4 in neutrophil activation and acute lung injury. *Am J Respir Cell Mol Biol*. 2016;54(3):370–83.
295. Hamanaka K, Jian M-Y, Townsley MJ, King JA, Liedtke W, Weber DS, et al. TRPV4 channels augment macrophage activation and ventilator-induced lung injury. *Am J Physiol-Lung Cell Mol Physiol*. 2010;299(3):L353–62.
296. Vassilieva IO, Negulyaev YA, Marakhova II, Semenova SB. TRPV5 and TRPV6 calcium channels in human T cells. *Cell Tissue Biol*. 2008;2(6):584–9.

297. Vassilieva IO, Tomilin VN, Marakhova II, Shatrova AN, Negulyaev YA, Semenova SB. Expression of transient receptor potential vanilloid channels TRPV5 and TRPV6 in human blood lymphocytes and jurkat leukemia T cells. *J Membr Biol*. 2013;246(2):131–40.
298. Michalick L, Kuebler WM. TRPV4—a missing link between mechanosensation and immunity. *Front Immunol*. 2020. <https://doi.org/10.3389/fimmu.2020.00413/full>.
299. Hartmannsgruber V, Heyken W-T, Kacic M, Kaistha A, Grgic I, Harteneck C, et al. Arterial response to shear stress critically depends on endothelial TRPV4 expression. *PLoS ONE*. 2007;2(9):e827.
300. Baratchi S, Knoerzer M, Khoshmanesh K, Mitchell A, McIntyre P. Shear stress regulates TRPV4 channel clustering and translocation from adherens junctions to the basal membrane. *Sci Rep*. 2017;7(1):15942.
301. Swain SM, Liddle RA. Shear stress-induced pathological changes in endothelial cells occur through Piezo1 activation of TRPV4. 2020.
302. Mendoza SA, Fang J, Guterman DD, Wilcox DA, Bubolz AH, Li R, et al. TRPV4-mediated endothelial Ca²⁺ influx and vasodilation in response to shear stress. *Am J Physiol Heart Circ Physiol*. 2010;298(2):H466–76.
303. Rajasekhar P, Poole DP, Veldhuis NA. Role of nonneuronal TRPV4 signaling in inflammatory processes. *Adv Pharmacol*. 2017;79:117–39.
304. Thodeti CK, Matthews B, Ravi A, Mammoto A, Ghosh K, Bracha AL, Ingber DE. TRPV4 channels mediate cyclic strain-induced endothelial cell reorientation through integrin-to-integrin signaling. *Circ Res*. 2009;104(9):1123–30.
305. Song X, Sun Z, Chen G, Shang P, You G, Zhao J, et al. Matrix stiffening induces endothelial dysfunction via the TRPV4/microRNA-6740/endothelin-1 mechanotransduction pathway. *Acta Biomater*. 2019;100:52–60.
306. Dutta B, Goswami R, Rahaman SO. TRPV4 plays a role in matrix stiffness-induced macrophage polarization. *Front Immunol*. 2020;11:570195.
307. Inoue R, Jian Z, Kawarabayashi Y. Mechanosensitive TRP channels in cardiovascular pathophysiology. *Pharmacol Ther*. 2009;123(3):371–85.
308. Narita K, Sasamoto S, Koizumi S, Okazaki S, Nakamura H, Inoue T, Takeda S. TRPV4 regulates the integrity of the blood-cerebrospinal fluid barrier and modulates transepithelial protein transport. *FASEB J*. 2015;29(6):2247–59.
309. Rosenkranz SC, Shaposhnikov A, Schnappauff O, Epping L, Vieira V, Heidermann K, et al. TRPV4-mediated regulation of the blood brain barrier is abolished during inflammation. *Front Cell Dev Biol*. 2020;8:849.
310. Potla R, Hirano-Kobayashi M, Wu H, Chen H, Mammoto A, Matthews BD, Ingber DE. Molecular mapping of transmembrane mechanotransduction through the β 1 integrin-CD98hc-TRPV4 axis. *J Cell Sci*. 2020. <https://doi.org/10.1242/jcs.248823>.
311. Strotmann R, Schultz G, Plant TD. Ca²⁺-dependent potentiation of the nonselective cation channel TRPV4 is mediated by a c-terminal calmodulin binding site. *J Biol Chem*. 2003;278(29):26541–9.
312. Zhao H, Zhang K, Tang R, Meng H, Zou Y, Wu P, et al. TRPV4 blockade preserves the blood-brain barrier by inhibiting stress fiber formation in a rat model of intracerebral hemorrhage. *Front Mol Neurosci*. 2018. <https://doi.org/10.3389/fnmol.2018.00097>.
313. Fabian A, Bertrand J, Lindemann O, Pap T, Schwab A. Transient receptor potential canonical channel 1 impacts on mechanosignaling during cell migration. *Pflügers Arch Eur J Physiol*. 2012;464(6):623–30.
314. Eijkelkamp N, Quick K, Wood JN. Transient receptor potential channels and mechanosensation. *Annu Rev Neurosci*. 2013;36:519–46.
315. Yao X, Garland CJ. Recent developments in vascular endothelial cell transient receptor potential channels. *Circ Res*. 2005;97(9):853–63.
316. Hicks K, O'Neil RG, Dubinsky WS, Brown RC. TRPC-mediated actin-myosin contraction is critical for BBB disruption following hypoxic stress. *Am J Physiol Cell Physiol*. 2010;298(6):C1583–93.
317. Trebak M, Lemonnier L, Smyth JT, Vazquez G, Putney JW Jr. Phospholipase C-coupled receptors and activation of TRPC channels. *Handb Exp Pharmacol*. 2007;179:593–614.
318. Ningoo M, Plant LD, Greka A, Logothetis DE. PIP(2) regulation of TRPC5 channel activation and desensitization. *J Biol Chem*. 2021;296:100726.
319. Adebisi A, Thomas-Gatewood CM, Leo MD, Kidd MW, Neeb ZP, Jaggar JH. An elevation in physical coupling of type 1 inositol 1,4,5-trisphosphate (IP 3) receptors to transient receptor potential 3 (TRPC3) channels constricts mesenteric arteries in genetic hypertension. *Hypertension*. 2012;60(5):1213–9.
320. Freichel M, Vennekens R, Olausson J, Stolz S, Philipp SE, Weissgerber P, Flockerzi V. Functional role of TRPC proteins in native systems: implications from knockout and knock-down studies. *J Physiol*. 2005;567(Pt 1):59–66.
321. Singh I, Knezevic N, Ahmmed GU, Kini V, Malik AB, Mehta D. Gq-TRPC6-mediated Ca²⁺ entry induces RhoA activation and resultant endothelial cell shape change in response to thrombin. *J Biol Chem*. 2007;282(11):7833–43.
322. Thilo F, Vorderwülbecke BJ, Marki A, Krueger K, Liu Y, Baumunk D, et al. Pulsatile atheroprone shear stress affects the expression of transient receptor potential channels in human endothelial cells. *Hypertension*. 2012;59(6):1232–40.
323. Du J, Fu J, Xia XM, Shen B. The functions of TRPP2 in the vascular system. *Acta Pharmacol Sin*. 2016;37(1):13–8.
324. Qu Y-Y, Wang L-M, Zhong H, Liu Y-M, Tang N, Zhu L-P, et al. TRPC1 stimulates calcium-sensing receptor-induced store-operated Ca²⁺ entry and nitric oxide production in endothelial cells. *Mol Med Rep*. 2017;16(4):4613–9.
325. Sharif-Naeini R, Folgering JH, Bichet D, Duprat F, Lauritzen I, Arhatte M, et al. Polycystin-1 and -2 dosage regulates pressure sensing. *Cell*. 2009;139(3):587–96.
326. Aboulaiwi WA, Takahashi M, Mell BR, Jones TJ, Ratnam S, Kolb RJ, Nauli SM. Ciliary polycystin-2 is a mechanosensitive calcium channel involved in nitric oxide signaling cascades. *Circ Res*. 2009;104(7):860–9.
327. Cha S-K, Kim J-H, Huang C-L. Flow-induced activation of TRPV5 and TRPV6 channels stimulates Ca(2+)-activated K(+) channel causing membrane hyperpolarization. *Biochem Biophys Acta*. 2013;1833(12):3046–53.
328. Shuvaev A, Kuvacheva NV, Morgun A, Khilazheva E, Salmina AB. The role of ion channels expressed in cerebral endothelial cells in the functional integrity of the blood-brain barrier (review). *Sovremennye tehnologii v medicine*. 2016;8:241–50.
329. Yankaskas CL, Bera K, Stoletoev K, Serra SA, Carrillo-Garcia J, Tuntithavornwat S, et al. The fluid shear stress sensor TRPM7 regulates tumor cell intravasation. *Sci Adv*. 2021;7(28):3457.
330. Numata T, Shimizu T, Okada Y. Direct mechano-stress sensitivity of TRPM7 channel. *Cell Physiol Biochem*. 2007;19(1–4):1–8.
331. Liu Y-S, Liu Y-A, Huang C-J, Yen M-H, Tseng C-T, Chien S, Lee OK. Mechanosensitive TRPM7 mediates shear stress and modulates osteogenic differentiation of mesenchymal stromal cells through osterix pathway. *Sci Rep*. 2015;5(1):16522.
332. Alvarado MG, Thakore P, Earley S. Transient receptor potential channel ankyrin 1: a unique regulator of vascular function. *Cells*. 2021. <https://doi.org/10.3390/cells10051167>.
333. Moparthi L, Zygmunt PM. Human TRPA1 is an inherently mechanosensitive bilayer-gated ion channel. *Cell Calcium*. 2020;91:102255.
334. Rostami A, Ciric B. Role of Th17 cells in the pathogenesis of CNS inflammatory demyelination. *J Neurol Sci*. 2013;333(1–2):76–87.
335. Rudolph H, Klopstein A, Gruber I, Blatti C, Lyck R, Engelhardt B. Postarrest stalling rather than crawling favors CD8 + over CD4 + T-cell migration across the blood–brain barrier under flow in vitro. *Eur J Immunol*. 2016;46(9):2187–203.
336. Abadier M, Pramod AB, McArdle S, Marki A, Fan Z, Gutierrez E, et al. Effector and regulatory T cells roll at high shear stress by inducible tether and sling formation. *Cell Rep*. 2017;21(13):3885–99.
337. Sundd P, Gutierrez E, Koltsova EK, Kuwano Y, Fukuda S, Pospieszalska MK, et al. 'Slings' enable neutrophil rolling at high shear. *Nature*. 2012;488(7411):399–403.
338. Dusi S, Angiari S, Pietronigro EC, Lopez N, Angelini G, Zenaro E, et al. LFA-1 controls Th1 and Th17 motility behavior in the inflamed central nervous system. *Front Immunol*. 2019;10:2436.
339. Woodfin A, Voisin M-B, Nourshargh S. PECAM-1: a multi-functional molecule in inflammation and vascular biology. *Arterioscler Thromb Vasc Biol*. 2007;27(12):2514–23.
340. Kalinowska A, Losy J. PECAM-1, a key player in neuroinflammation. *Eur J Neurol*. 2006;13(12):1284–90.
341. Chien S. Effects of disturbed flow on endothelial cells. *Ann Biomed Eng*. 2008;36(4):554–62.
342. Alexander JS, Alexander BC, Eppihimer LA, Goodyear N, Haque R, Davis CP, et al. Inflammatory mediators induce sequestration of VE-cadherin in cultured human endothelial cells. *Inflammation*. 2000;24(2):99–113.

343. Koivisto A-P, Belvisi MG, Gaudet R, Szallasi A. Advances in TRP channel drug discovery: from target validation to clinical studies. *Nature Reviews Drug Discovery*. 2021.
344. Miller F, Björnsson M, Svensson O, Karlsten R. Experiences with an adaptive design for a dose-finding study in patients with osteoarthritis. *Contemp Clin Trials*. 2014;37(2):189–99.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.